Edited by Erasmo RECAMI

TACHYONS, Monopoles, and Related Topics

North-Holland

TACHYONS, MONOPOLES, AND RELATED TOPICS

Proceedings of the First Session of the Interdisciplinary Seminars on "Tachyons and Related Topics"

Erice, 1–15 September, 1976

Edited by:

ERASMO RECAMI

Institute of Theoretical Physics,
University of Catania, Italy,
and
I.N.F.N. Catania

"Ettore Majorana" International Centre for Scientific Culture
ANTONINO ZICHICHI, *Director*

Interdisciplinary Seminars
EUGENE P. WIGNER, *Director*

First Session: Tachyons and Related Topics
ERASMO RECAMI, *Director*

1978

NORTH-HOLLAND PUBLISHING COMPANY – AMSTERDAM • NEW YORK • OXFORD

© North-Holland Publishing Company, 1978

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright owner.

ISBN: 0 444 85165 8

Publishers

NORTH-HOLLAND PUBLISHING COMPANY
AMSTERDAM • NEW YORK • OXFORD

Sole distributors for the U.S.A. and Canada
ELSEVIER NORTH-HOLLAND, INC.
52 VANDERBILT AVENUE,
NEW YORK, N.Y. 10017

Library of Congress Cataloging in Publication Data

Main entry under title:

Tachyons, monopoles, and related topics.

Held under the auspices of Ettore Majorana International Centre for Scientific Culture.
Includes index.
1. Tachyons--Congresses. 2. Quantum field theory--Congresses. I. Recami, Erasmo. II. Ettore Majorana International Centre for Scientific Culture.
QC793.5.T322T3 539.7'21 78-5803
ISBN 0-444-85165-8

PRINTED IN THE NETHERLANDS

C O N T E N T S

PREFACE v

P A R T A: T A C H Y O N S

PART A1: TACHYONS IN RELATIVISTIC CLASSICAL THEORIES:

E.RECAMI: AN INTRODUCTORY VIEW ABOUT SUPERLUMINAL FRAMES AND TACHYONS 3
1. Rivisiting the Postulates of Special Relativity (SR)
2. Causality in SR
3. Some consequences
4. Extended Relativity: Historical Remarks
5. Preliminaries (and caveats) on Tachyons
6. The Generalized Lorentz Transformations (GLT)
7. Tachyons and Causality
8. Classical Physics for Tachyons
9. Tachyons and Black-holes
10. Virtual Particles and Tachyons
11. Astrophysics and Superluminal Objects
12. Note Added

J.S.TREFIL: TACHYONS AND THE CAUSAL PARADOX 27

H.C.CORBEN: ELECTROMAGNETIC AND HADRONIC PROPERTIES OF TACHYONS 31
1. Attitudes About Tachyons
2. Superluminal Trasformations
3. Electromagnetic Properties of Tachyons
4. Tachyons in Quantum Mechanics
5. A relativistically invariant hadron Bootstrap

E.C.G.SUDARSHAN: TACHYONS AND THE SEARCH FOR A PREFERRED FRAME 43
1. Bradyons, Luxons and Tachyons
2. Transcendent Tachyons and the Reinterpretation Principle
3. The dazzling Tachyon Sky
4. Search for a Preferred Frame
5. Concluding remarks

Ya.P.TERLETSKY: ABOUT TACHYONS 47
1. The Electromagnetic Radiation of a Charged Tachyon
2. Some properties of Tachyons and Causality

K.Tahir SHAH: TACHYONS AND THEORY OF QUASI-CATASTROPHES 49

A.J.KÁLNAY: COMPLEX PHYSICAL QUANTITIES AND SPACE-LIKE STATES 53
1. Introduction
2. Can Complex Numbers be the result of Experiments?
   2a - The case of classical physics
   2b - Quantum mechanics and non-normal operators
   2c - Q.M., non-normal operators and extended variables
   2d - Levy-Leblond's approach
3. On Tachyon Physics

H.HÄNNI and E.HUGENTOBLER: TACHYONS AND EXPERIMENTAL PHYSICS 61
1. General Remarks
2. Two possible Tachyon Detectors
2a - A Cherenkov detector for tachyons
2b - A time-of-flight (TOF) tachyon detecting system

R.MIGNANI: QUATERNIONS AND LORENTZ TRANSFORMATIONS 67
1. Introduction
2. Algebraic and metric properties of Quaternions
3. Minquats
4. Subluminal Lorentz Transformations
5. Superluminal Lorentz Transformations
6. Complex Minkowski space-time, Magnetic Monopoles and Tachyons

J.D.EDMONDS Jr.: COVARIANT QUANTUM EQUATIONS IN CURVED SPACE-TIME, LORENTZ COVARIANCE AND TACHYONS 79
1. Hypercomplex numbers, Mass and Space-Time
2. Group structures in Nature's number system
3. Natural quantum equations
4. Tachyons and Absolute Motion
5. Curved Space Derivatives
6. Some considerations on the Gravity curvature equation

M.CAMENZIND: TACHYON MATTER 89
1. Introduction: The standard picture of the particle world
2. Spacelike representations of the Poincaré Group
3. Some properties of the Mass Shell
4. The One-Particle tachyon states
5. Tachyonic Matter and Curved Space-Times
6. Tachyonic Matter in the Gravitational Collapse

V.DE SABBATA: BLACK HOLES AND MOTION OF TEST PARTICLES 99

M.PAVŠIČ: UNIFIED DESCRIPTION OF BRADYONS AND TACHYONS BASED ON THE DYNAMICAL SO(4,2) SYMMETRY 105
1. Introduction
2. The law of motion in the flat (η^a)-space
3. Motion of a photon
4. The law of motion in a curved (η^a)-space
5. Relation between the *scale* κ and the type of particle

G.ARCIDIACONO: TACHYONS AND MAGNETIC MONOPOLES IN THE DE SITTER UNIVERSE 115
1. Introduction
2. The De Sitter-Castelnuovo space-time
3. The transformations of the Fantappié Group
4. The Cosmical Expansion and the red-shift laws
5. The De Sitter Universe and Mechanics
6. Magnetic Monopoles and Plasma's Physics
7. Magnetic Monopoles, hyperdense matter and Cosmology

R.GOLDONI: FASTER-THAN-LIGHT PHENOMENA IN SPECIAL AND IN GENERAL RELATIVITY 125
1. Introduction
2. Special Relativity
3. Quantum field theory

61 4. General Relativity
5. Black-and-White-Holes
6. Conclusions

PART A2: TACHYONS IN RELATIVISTIC QUANTUM THEORIES

67 A.O.BARUT: SPACE-LIKE STATES IN RELATIVISTIC QUANTUM THEORY 143
1. Introduction
2. Space-like Representations of the Poincaré Group II
2a - Formulas for time-like case
2b - Formulas for space-like case
2c - Induced Representation from Little Groups
yons 3. Localization of Space-like States
3a - The Imprimitivity System
79 3b - Covariant formulation
3c - New forms of localizability for light-like and space-like cases
4. Passage to Covariant Wave Equations
4a - Induced representations and wave equations
4b - Induction from a Dynamical Group G
5. Physical Interpretation of Space-like Solutions
5a - Passage to space-time
89 5b - Relativistic Dynamics
5c - Transformations connecting inequivalent representations
6. Space-like States in S-matrix theory and in Infinite-component Wave Equations

K.KAMOI and S.KAMEFUCHI: TACHYONS AS VIEWED FROM QUANTUM FIELD THEORY 159
1. Introduction
2. Invariant Delta Functions
99 3. Canonical Quantization
4. An Alternative Approach
105 5. Additional Remarks

H.B.NIELSEN: TACHYONS IN FIELD THEORY: CONVENTIONAL TREATMENT 169

E.VAN DER SPUY: TACHYONS AND VIRTUAL FIELDS FOR ELEMENTARY PARTICLES IN STRONG INTERACTIONS 175
I - TACHYONS: WHY? 175
1. Introduction
ERSE 115 2. The Reggeization of the Field
3. Unstable Compounds: Resonances and Complex Regge Poles
II - TACHYONS: HOW? 183

D.SHAY and K.L.MILLER: SPIN-S TACHYONS AND THE PROPAGATION OF TACHYON WAVES 185
D.SHAY (I part): SPIN-S TACHYONS 185
1. Introduction
2. Spinors in the Zero Energy Frame
3. Transformation of Tachyon Spinors
125 4. Invariant Integral
5. Conclusions
D.SHAY and K.L.MILLER: PROPAGATION OF TACHYON WAVES 189
1. Introduction

2. Phase and Group Velocity
3. Wave Packets
4. Delta Function Pulses
5. Conclusions

J.KRÜGER: JOINT DISTRIBUTIONS AND TACHYONS 195
1. Introduction
2. Wigner-Distribution and its Generalization in the Non-relativistic Theory
3. Relativistic J.D.F. and Green Functions

PART B:

MONOPOLES, TACHYONS, AND RELATED TOPICS

E.FERRARI: FORMULATION OF ELECTRODYNAMICS WITH MAGNETIC MONOPOLES 203
1. Introduction
2. The Basic Features of the theory of Maxwell-Dirac Monopoles
3. The first Dirac paper on Monopoles
4. The Dirac Formulation of Electrodynamics with Monopoles
5. The Schwinger Formulation
6. The Zwanziger Formulation
7. Feynman Diagrams
8. The Cabibbo-Ferrari Formulation
9. The Wu-Yang Approach
10. Conclusions
11. Note Added

A.O.BARUT: ELECTRODYNAMICS WITH STRINGS AND MONOPOLES 227
1. Introduction
2. The Completion of Dirac's Action Principle and Removal of Difficulties
3. Coordinate Independent Formulation on Manifolds and Topological Aspects
4. Some new Quantum Effects: Parity and Superselection Rules
5. Realistic Hadron Models: Dyonium - Extended Dyonium

G.PARISI: NON CONVENTIONAL MONOPOLES IN FIELD THEORY 233

M.GARCÍA-SUCRE: REFERENCE FRAMES AND THE VELOCITY OF LIGHT 235

R.CASALBUONI: COLORED QUARKS AND OCTONIONS 247
1. Colour
2. Octonions in Mathematics
3. The Description of the Physical States
4. Octonions and Exceptional Lie Groups

G.ZIINO: CHIRALITY INVARIANCE AND "CHIRAL" FIELDS 261

APPENDIX

E.RECAMI: NEW EVIDENCE ON THE LIFE OF ETTORE MAJORANA 269

1. Foreword
2. The last letters of Ettore Majorana
3. The Man and the Scientist

LIST OF PARTICIPANTS 279

AUTHOR INDEX 283

195

stem,
ding

PREFACE

This volume contains the Proceedings of the Erice First Session of the Interdisciplinary Seminars on <<Tachyons and Related Topics>>. It is surely the first book (in the world) that deals with the extension of the Relativistic (Classical and Quantal) Theories to faster-than-light frames and/or objects, and with the application of such extended theories to a better comprehension (also) of the ordinary physical world.

All contributions, initially expounded in lectures (and seminars) at the Erice International Meeting, have been up-to-dated up to Jan., 1978. The list of the authoritative lecturers and of their themes (showing a large spectrum of topics) is given in the Contents. The high standard of the contributions should appear from the texts. Let us only add that this volume is likely to contain many permanent results.

Due to the possible connections between Super-luminal "charges" and "monopoles", and to the possible rôle of Tachyons in hadron-structure and elementary particle interactions, large space is also devoted to MONOPOLES in themselves, as well as to hadron constituents in themselves (for these topics see e.g. the chapters by A.O.Barut, R.Casalbuoni, E.Ferrari, H.B.Nielsen, G.Parisi, etc.).

Research about Tachyons, in particular, has been constantly growing up during the last ten years and the related scientific literature already consists of many hundreds of papers. This book hopes to become an initial "reference point" for facilitating theoretical and experimental research on tachyons and related topics, and to offer a vehicle towards the understanding of such kinds of problems to the interested scientists and to the university teachers of relativistic theories. The subjects range from the solution of the Causal "paradoxes" to the application of tachyon "philosophy" to Astrophysics. Namely —with regard to Tachyons— these Proceedings deal e.g. with: Tachyons in Special Relativity, in General Relativity, in Projective Relativity and in Conformal Relativity; Space-like states from the group theoretical viewpoint; Quantum Field Theory of tachyons; Causality; External and internal symmetries in sub-nuclear physics; Tachyons' spin and "localization"; Black-holes and Tachyons; Astrophysics, cosmology and Superluminal objects; conventional and non-conventional Monopoles, and tachyons; Complex physical quantities, the nature of space-time and space-like states; Structure of the vacuum and tachyons; Virtual Particles and Tachyons in Resonance formation and elementary particle physics; Usefulness of the Quaternion and Octonion formalisms and of Catastrophes theory in building up relativistic physics; Electromagnetic properties of tachyons; Joint distributions and tachyons; and so on. The experimental question is not forgotten. Let us notice, at this point, that the present experimental situation might often be criticized, since most experiments till now performed looking for tachyons seem devoid of a careful theoretical ground.

The structure of this book has been conceived as follows: we start with the problem of extending the Classical (relativistic) theories, by considering firstly the Special Relativity and secondly the General Relativity; then we pass to the extension of Relativistic Quantum theories; and finally we discuss Monopoles and topics related to tachyons and/or monopoles. A paper containing various arguments can appear twice in the Contents. Even if the matter of the present volume is rather recent, nevertheless many chapters here presented show a substantial,

mutual agreement, complementing one another. However, different lines of thought can be met; for instance, Goldoni's approach is partially different from the Introductory one.

In the Appendix, new biographical evidence is reported about the figure of Ettore MAJORANA (the brilliant physicist who mysteriously disappeared at 31 in 1938), whom the Erice "Centre" is dedicated to; such an Appendix originates from an evening talk organized at the kind invitation of the Director of the parallel course, held at Erice during the same period. The book ends with the List of Participants and the Author Index (the latter being due to Dr.L.R.Baldini).

Two defects of this volume are the absence of a contribution from Prof.G.Bernardini, announced as one of the lecturers — who unfortunately could not join us due to a personal hitch — and the absence of the text of the beautiful survey on the topics discussed in the parallel course (about Nuclear Physics), kindly delivered to us by Sir Denys Wilkinson, whose courtesy is here gratefully acknowledged.

Thanks are due to Dr.R.Mignani, who has been the Scientific Secretary for our Erice Conference.

This Erice Meeting was mainly supported by I.N.F.N. (Prof.A.Gigli, the President), with a small contribution from the Università di Catania, Catania, Italy. The preliminary work for the preparation of the present Proceedings is expected to be supported in part by C.N.R. The editor, moreover, acknowledges the hospitality received at the Niels Bohr Institut, Copenhagen, Denmark, from where part of the Conference was organized, and at the Astrophysics Department, University of Oxford, U.K., where part of this Proceedings-volume was prepared; thanks are due to Profs.A.Bohr, H.B.Nielsen, P.Olesen and to Profs.K.W.Allen, D.E.Blackwell, D.W.Sciama, respectively.

I wish to thank Profs. A.Agodi, I.Agostini, Mrs.Baldini R., A.O.Barut, G.Bernardini, P.Caldirola, F.Catara, H.C.Corben, V.De Sabbata, J.Deutsch, H.D.Engelhardt, Mrs.M.Fallon, A.Galli, A.Gigli B., A.J.Kálnay, E.Oliveri, G.Rodolico, R.Ruffini, E.P.Wigner, D.Wilkinson, S.A.Wouthuysen and A.Zichichi for their kind, various interest; Dr.A.Gabriele and Miss P.Savalli (Majorana Center) for their meeting-organization assistance in Erice; and all the Participants of our Meeting for their competent, courteous contribution and their affectionate, warm presence.

Finally, many thanks are sincerely due to the Senior Physics Editor Dr.P.S.H. Bolman (and Miss A.Vickerson), of the North Holland Publ.Co., for having decided (and taken care of) the birth of the present book.

Catania, Feb. 1978

E.R.

Proceedings of the

SIR ARTHUR EDDINGTON CENTENARY SYMPOSIUM

VOLUME 3

GRAVITATIONAL RADIATION AND RELATIVITY

Editors
J Weber
T M Karade

World Scientific

A MODEL-THEORY FOR TACHYONS IN TWO DIMENSIONS (°)

Erasmo Recami (*) and Waldyr A.Rodrigues(")

Istituto Nazionale di Fisica Nucleare, Sezione di Catania, Catania, Italy

Abstract - The subject of *Tachyons*, even if still speculative, may deserve some attention for reasons that can be divided into a few categories, two of which we want preliminarily to mention right now: (i) the larger scheme that one tries to build up in order to incorporate space-like objects in the relativistic theories can allow a better understanding of many aspects of the *ordinary* relativistic physics, even if Tachyons would not exist in our cosmos as "asymptotically free" objects; (ii) Superluminal classical objects can have a role in elementary particle interactions (and perhaps even in astrophysics); and it might be tempting to verify how far one can go in reproducing the quantum-like behaviour at a classical level just by taking account of the possible existence of faster-than-light classical particles.

This article is divided in two parts, the first one having nothing to do with tachyons. In fact, to prepare the ground, in Part I (Sect.2) we shall merely show that Special Relativity —even *without* tachyons— can be given a form such to describe both particles and anti-particles. The plan of Part II is confined only to a "model-theory" of Tachyons in two dimensions, for the reasons stated in Sect.3.

(°) Supported in part by MPI, CNR, and by IBM-do-Brazil.

(*) Istituto Dip.le di Fisica, Università Statale di Catania, Catania, Italy. Present address: Dept. of Applied Mathem., State University at Campinas, Campinas, S.P., Brazil, on leave of absence from Catania, Italy.

(") Dept. of Applied Mathem., State University at Campinas, Campinas, S.P., Brazil.

1. INTRODUCTION

The subject of *Tachyons*, even if still speculative, may deserve some attention for reasons that can be divided into a few categories, two of which we want preliminarly to mention right now: (i) the larger scheme that one tries to build up in order to incorporate space-like objects in the relativistic theories can allow a better understanding of many aspects of the *ordinary* relativistic physics, even if Tachyons would not exist in our cosmos as "asymptotically free" objects; (ii) Superluminal classical objects can have a role in elementary particle interactions (and perhaps even in astrophysics); and it might be tempting to verify how far one can go in reproducing the quantum-like behaviour at a classical level just by taking account of the possible existence of faster-than-light classical particles.

This article is divided in two parts, the first one having nothing to do with tachyons. In fact, to prepare the ground, in Part I (Sect. 2) we shall merely show that Special – even *without* tachyons – can be given a form such to describe both particles and anti-particles. The plan of Part II is confined only to a "model-theory" of Tachyons in two dimensions, for the reasons stated in Sect. 3.

PART I : PARTICLES AND ANTIPARTICLES IN SPECIAL RELATIVITY (SR)

2. SPECIAL RELATIVITY WITH ORTHO- AND ANTI-CHRONOUS LORENTZ TRANSFORMATIONS.

In this Part I we shall *forget* about Tachyons.

From the ordinary postulates of Special Relativity (SR) it follows that in such a theory – which refers to the class of Mechanical and Electromagnetic Phenomena – the class of reference-frames equivalent to a given inertial frame is obtained by means of transformations L (Lorentz Transformations, LT) which satisfy the following sufficient requirements: (i) to be linear;

$$x'^{\mu} = L^{\mu}_{\nu}\, x^{\nu} \ ; \tag{1}$$

(ii) to preserve space-isotropy (with respect to electromagnetic and mechanical phenomena); (iii) to form a group; (iv) to leave the quadratic form invariant:

$$\eta_{\mu\nu}\, dx^{\mu} dx^{\nu} = \eta'_{\alpha\beta}\, dx'^{\alpha} dx'^{\beta} \ . \tag{2}$$

From condition (i), if we confine ourselves to sub-luminal speeds, it follows that in eq.(2):

$$\eta'_{\alpha\beta} = \mathrm{diag}(+1, -1, -1, -1) = \eta_{\mu\nu}. \tag{3}$$

Eqs. (1)-(3) imply that $\det L^2 = 1$; $(L^{o}_{o})^2 \geq 1$. The set of all sub-luminal (Lorentz) transformations satisfying all our conditions consists — as is well-known — of four pieces, which form a noncompact, non-connected group (the Full Lorentz Group). Wishing to confine ourselves to space-time "rotations" only, i.e. to the case $\det L = +1$, we are left with the two pieces:

$$\begin{cases} \{L^{\uparrow}_{+}\} : L^{o}_{o} \geq +1 \quad ; \quad \det L = +1 \ ; & (4\underline{a}) \\ \{L^{\downarrow}_{+}\} : L^{o}_{o} \leq -1 \quad ; \quad \det L = -1 \ , & (4\underline{b}) \end{cases}$$

which give origin to the group of the proper (orthochronous *and* antichronous) transformations

$$\mathcal{L}_{+} \equiv \mathcal{L}^{\uparrow}_{+} \cup \mathcal{L}^{\downarrow}_{+} \equiv \{L^{\uparrow}_{+}\} \cup \{L^{\downarrow}_{+}\} \tag{5}$$

and to the subgroup of the (ordinary) proper ortochronous transformations

$$\mathcal{L}^{\uparrow}_{+} \equiv \{L^{\uparrow}_{+}\} \ , \tag{6}$$

both of which being, incidentally, invariant subgroups of the Full Lorentz Group. For reasons to be seen later on, let us rewrite $\mathcal{L}_+$ as follows

$$\mathcal{L}_+ = \mathcal{L}_+^\uparrow \otimes \mathbf{Z}(2) \; ; \quad \mathbf{Z}(2) \equiv \{\sqrt[2]{+1}\} \equiv \{+1, -1\}. \tag{5'}$$

We shall skip in the following, for simplicity's sake, the subscript + in the transformations $L_+^\uparrow$, $L_+^\downarrow$. Given a transformation $\overline{L}^\downarrow$, another transformation $\overline{L}^\uparrow \in \mathcal{L}_+^\uparrow$ always exists such that

$$\overline{L}^\downarrow = (-\mathbb{1}) \cdot \overline{L}^\uparrow, \quad \forall \overline{L}^\downarrow \in \mathcal{L}_+^\downarrow \; , \tag{7}$$

and vice-versa. Such a one-to-one correspondence allows us to write formally

$$\mathcal{L}_+^\downarrow = -\mathcal{L}_+^\uparrow \; . \tag{7'}$$

It follows in particular that the central elements of $\mathcal{L}_+$ are: $\mathbb{C} \equiv \equiv (+\mathbb{1}, -\mathbb{1})$.

Usually, even the piece (4b) is discarded. Our present aim is to show — on the contrary — that a physical meaning can be attributed also to the transformations (4b). Confining ourselves here to the *active* point of view (cf. Recami and Rodrigues[1] and references therein), we wish precisely to show that the theory of SR, once based on the *whole* proper Lorentz group (5) and not only on its orthochronous part, will describe a Minkowski space-time populated by both matter and antimatter.

2.1. The Stückelberg-Feynman "switching principle" in SR.

Besides the usual postulates of SR (Principle of Relativity, and Light-Speed Invariance), let us assume — as commonly admitted, e.g. for the reasons in Garuccio *et al.*[9], Mignani and Recami[3] — the following:

Assumption - ((negative-energy objects travelling forward in time do *not* exist)) . We shall give this Assumption, later on, the status of a fundamental postulate.

Let us therefore start from a positive-energy particle P travelling forward in time. As wellknown, any orthochronous LT (4a) transforms it into another particle still endowed with positive energy and motion forward in time. On the contrary, any antichronous (=non-orthochronous) LT (4b) will change sign — among the others — to the time-components of *all the four-vectors* associated with P. Any $L^{\uparrow}$ will transform Γ into a particle P' endowed in particular with negative energy *and* motion backwards in time. (Fig. 1).



Fig. 1

In other words, SR together with the natural Assumption above *implies* that a particle going backwards in time (Gödel[4]) (Fig. 1) corresponds in the four-momentum space, Fig. 2, to a particle carrying negative energy; and, vice-versa, that changing the energy sign in one space corresponds to changing the sign of time in the dual space. It is then easy to see that these two paradoxical occurrences ("negative energy" and "motion backwards in time") give rise to a phenomenon that



Fig. 2

any observer will describe in a quite *orthodox* way, when they are — as they actually are — simultaneous (Recami[5,6] and refs. therein).

Notice, namely, that: (i) every observer (a macro-object) explores space-time, Fig. 1, in the positive t-direction, so that we shall meet B as the first and A as the last event; (ii) emission of positive quantity is equivalent to absorption of negative quantity, as (-) · (-) = = (+) · (+); and so on.

Let us now suppose (Fig. 3) that a particle P' with negative energy (and e.g. charge -e) moving backwards in time is emitted by A at time t_1 and absorbed by B at time $t_2 < t_1$. Then, it follows that at time t_1 the object A "looses" negative energy and charge, i. e. *gains* positive energy and charge. And that at time $t_2 < t_1$ the object B "gains" negative energy and charge, i.e. *looses* positive energy and charge. The physical phenomenon here described is nothing but the exchange *from* B *to* A of a particle Q with *positive* energy, charge +e, and going *forward* in time. Notice that Q has, however, charges *opposite* to P'; this means that in a sense the present "switching procedure" (previously called "RIP") effects a "charge conjugation" C, among



Fig. 3(a)

$[t_1 > t_2]$

(t_1, x_1) A $\xrightarrow{(P);-q;E<0;\overleftarrow{T};p<0}$ B (t_2, x_2)

$\xleftarrow{(Q);+q;E\geq 0;\vec{T};p>0}$

$[t<t']$

$$\text{ph} = \underset{(t,x)}{\boxed{A}} \xrightarrow[(+\lambda);\nu>0]{(+q);E>0;\vec{T};p>0} \underset{(t',x')}{\boxed{B}}$$

$$\hat{C}\hat{T}(\text{ph}) = \underset{(-t,x)}{\boxed{\bar{A}}} \xleftarrow[(+\lambda);\nu<0]{(-q);E>0;\vec{T};p<0} \underset{(-t',x')}{\boxed{\bar{B}}}$$

Fig.3(b)

$$\text{"}RIP\text{"}(\text{ph}) = \underset{(t,x)}{\boxed{A}} \xleftarrow[(+\lambda);\nu>0]{(-q);E<0;\overleftarrow{T};p<0} \underset{(t',x')}{\boxed{B}}$$

$$\hat{C}\hat{P}\hat{T}(\text{ph}) = \underset{}{\boxed{\bar{B}}} \xrightarrow[(-\lambda);\nu>0]{(-q);E>0;\vec{T};p>0} \underset{}{\boxed{\bar{A}}}$$

the others. Notice also that "charge", here and in the following, means *any* additive charge; so that our definitions of charge conjugation, etc., are more general than the ordinary ones (Recami and Mignani [7], hereafter called Review I; Recami [8]). Incidentally, such a switching

procedure has been shown to be equivalent to applying the chirality operation γ_5 (Recami and Ziino[9]).

2.2. Matter and Antimatter from SR

A close inspection shows the application of any antichronous transformation $L^{\downarrow}$, together with the switching procedure, to transform P into an object

$$Q \equiv \overline{P} \tag{8}$$

which is indeed the *antiparticle* of P. We are saying that *the concept of antimatter is a purely relativistic one*, and that, on the basis of the double sign in [c = 1]

$$E = \pm \sqrt{\vec{p}^2 + m_o^2} \quad , \tag{9}$$

the existence of antiparticles could have been predicted from 1905, exactly with the properties they actually exibited when later discovered, provided that recourse to the "switching procedure" had been made. We therefore maintain that the points of the lower hyperboloid sheet in Fig. 2 — since they corresponds not only to negative energy but also to motion backwards in time — represent the kinematical states of the *antiparticle* $\overline{P}$ (*of* the particle P represented by the upper hyperboloid sheet). Let us explicitly observe that the switching procedure exchanges the roles of *sources* and *detector*, so that (Fig. 1) any observer will describe B to be the source and A the detector of the antiparticle $\overline{P}$.

Let us stress that the switching procedure not only can, but *must* be performed, since any observer can do nothing but explore space-time along the positive time-direction. That procedure is merely the translation into a purely relativistic language of the Stückelberg-Feynman[10] "Switching principle". Together with our Assumption above, it can take the form of a "Third Postulate": ((Negative-energy objects

travelling forward in time do not exist; any negative-energy object P travelling backwards in time can and must be described as its anti-object $\bar{P}$ going the opposite way in space (but endowed with positive energy and motion forward in time)〉〉 . Cf. e.g. Caldirola and Recami,[11] Recami[6] and references therein.

2.3. Further remarks

a) Let us go back to Fig. 1. In SR, when based only on the two ordinary postulates, nothing prevents a priori the event A from influencing the event B. Just to forbid such a possibility we introduced our Assumption together with the Stückelberg-Feynman "Switching procedure". As a consequence, not only we eliminate any particle-motion backwards in time, but we also "predict" and naturally explain within SR the existence of antimatter.

b) The Third Postulate, moreover, helps solving the paradoxes connected with the fact that all relativistic equations admit, besides standard "retarded" solutions, also "advanced" solutions: The latter will simply represent antiparticles travelling the opposite way (Mignani and Recami[12]). For instance, if Maxwell equations admit solutions in terms of *outgoing* (polarized) photons of helicity $\lambda = +1$, then they will admit also solutions in terms of incoming (polarized) photons of helicity $\lambda = -1$; the actual intervention of one or the other solution in a physical problem depending only on the initial conditions.

c) Eqs. (7), (8) tell us that, in the case considered, any $\bar{L}^{\downarrow}$ has the same kinematical effect than its "dual" transformation $\bar{L}^{\uparrow}$ just defined through eq. (7), except for the fact that it moreover transforms P into its antiparticle $\bar{P}$. Eqs. (7), (7') then lead (Mignani and Recami[13,15]) to write

$$-\mathbb{1} \equiv \bar{P}\,\bar{T} = CPT \ , \qquad (10)$$

where the symmetry operations $\overline{P}$, $\overline{T}$ are to be understood in the "strong sense": For instance, $\overline{T}$ = reversal of the time-components of all fourvectors associated with the considered phenomenon (namely, inversion of the time *and* energy axes). We shall come back to this point. The discrete operations P,T have the ordinary meaning. When the particle considered in the beginning can be regarded as an extended object, Pavšič and Recami[16] have shown the "strong" operations $\overline{P}$, $\overline{T}$ to be equivalent to the space, time reflections acting on the space-time *both* external *and* internal to the particle world-tube.

Once accepted eq. (10), then eq.(7') can be written

$$\mathcal{L}_+^\downarrow = (\overline{P}\,\overline{T})\,\mathcal{L}_+^\uparrow \equiv (CPT)\mathcal{L}_+^\uparrow \ . \qquad (7'')$$

In particular, the total-inversion $\overline{L}^\downarrow = -\mathbf{1}$ transforms the process $a + b \to c + d$ into the process $\overline{d} + \overline{c} \to \overline{b} + \overline{a}$ without any change in the velocities.

d) All the ordinary relativistic laws (of Mechanics and Electromagnetism) are actually already covariant under the *whole* proper group $\mathcal{L}_+$, eq.(5), since they are CPT-symmetric besides being covariant under $\mathcal{L}_+^\uparrow$.

e) A few quantities that happened (cf. Sect. 5.17 in the following) to be Lorentz-invariant under the transformations $L^\uparrow \in \mathcal{L}_+^\uparrow$, are no more invariant under the transformations $L \in \mathcal{L}_+$. We have already seen this to be true for the *sign* of the additive charges, e.g. for the sign of the electric charge $\underline{e}$ of a particle P. The ordinary derivation of the electric-charge invariance is obtained by evaluating the integral flux of a current through a surface which, under $L^\uparrow$, does move, changing the angle formed with the current. Under $L^\downarrow \in \mathcal{L}_+^\downarrow$ the surface "rotates" so much with respect to the current (cf. also Figs. 6,12 in the following) that the current enters is through the opposite face; as a consequence, the integrated flux (i.e. the charge) changes sign.

PART II : BRADYONS AND TACHYONS IN SR

3. PRELIMINARIES ABOUT TACHYONS

Let us now take on the issue of Tachyons (T).

Tachyons, or space-like particles, are already known to exist as *internal, intermediate states* or exchanged objects. Can they also exist as "asymptotically free" objects?

We shall see that the particular — and unreplaceable — role in SR of the light-speed c in vacuum is due to its *invariance* (namely, to the experimental fact that c does not depend on the velocity of the source), and *not* to its being or not the maximal speed (Recami and Modica[17]).

Since a priori we know nothing about Ts, the safest way to build up a theory for them is trying to generalize the ordinary theories (starting with the classical relativistic one, only later on passing to the quantum field theory) through "minimal extensions", i.e. by performing modifications as small as possible. Only afer possessing a theoretical model we shall be able to start experiments: Let us remember that, not only good experiments are required before getting sensible ideas (Galilei[18,19]), but also a good theoretical background is required before sensible experiments can be performed.

The first step consists therefore in facing the problem of extending SR to Tachyons. In so doing, some authors limited themselves to consider objects both subluminal and Superluminal, always referred however to subluminal observers ("weak approach"). Other authors attempted on the contrary to generalize SR by introducing both subluminal observers (s) and Superluminal observers (S), and then by extending the Principle of Relativity ("strong approach"). This second approach is theoretically more worth of consideration (tachyons, e.g., get *real* proper-masses), but is meets of course the greatest obstacles. In fact, the extension of the Relativity Principle to Superluminal inertial frames seems to be straighforward only in the pseudo-Euclidean space-times M(n,n) having the same number n of space-axes and of time-axes.

For instance, when facing the problem of generalizing the Lorentz transformations to Superluminal frames in four dimensions one meets no-go theorems as Gorini's *et al.* (Gorini[20] and refs. therein), stating *no* such extensions exist which satisfy *all* the following properties: (i) to refer to the four-dimensional Minskowski space-time $M_4 \equiv M(1,3)$; (ii) to be real; (iii) to be linear; (iv) to preserve the space isotropy; (v) to preserve the light-speed invariance; (vi) to possess the prescribed group-theoretical properties.

We shall therefore start by sketching the simple, instructive and very promising "model-theory" in two dimensions (n=1).

Let us first revisit, however, the postulates of the ordinary SR.

4. THE POSTULATES OF SR REVISITED

Let us adhere to the ordinary postulates of SR. A suitable choice of Postulates is the following one (Review I; Maccarrone and Recami[21,22] and refs. therein):

1) First Postulate - Principle of Relativity: ⟨⟨The physical laws of Electromagnetism and Mechanics are covariant (= invariant in form) when going from an inertial frame $\underline{f}$ to another frame moving with constant velocity $\vec{u}$ relative to $\underline{f}$⟩⟩.

2) Second Postulate - "Space and time are homogeneous and space is isotropic". For future convenience, let us give this Postulate the form: ⟨⟨The space-time accessible to any inertial observer is four-dimensional. To each inertial observer the 3-dimensional Space appears as *homogeneous and isotropic*, and the 1-dimensional Time appears as *homogeneous*⟩⟩.

3) Third Postulate - Principle of Retarded Causality: ⟨⟨Positive-energy objects travelling backwards in time *do not exist*; and any negative-energy particle P travelling backwards in time can and must be described as its antiparticle $\overline{P}$, endowed with positive energy and motion forward in time (but going the opposite way in space)⟩⟩. See Sects. 2.1, 2.2.

The *First Postulate* is inspired to the consideration that all inertial frames should be *equivalent* (for a careful definition of "equivalence" see e.g. Recami[6]); notice that this Postulate does *not* impose any constraint on the relative speed $u \equiv |\vec{u}|$ of the two inertial observers, so that a priori $-\infty < u < +\infty$. The *Second Postulate* is justified by the fact that from it the conservation laws of energy, momentum and angular-momentum follow, which are well verified by experience (at least in our "local" space-time region); let us add the following comments: (i) The words homogeneous, isotropic refer to space-time properties assumed — as always — with respect to the electromagnetic and mechanical phenomena; (ii) Such properties of space-time are supposed by this Postulate to be covariant within the class of the inertial frames; this means that SR assumes the vacuum (i.e. space) to be "at rest" with respect to *every* inertial frame. The *Third Postulate* is inspired to the requirement that *for each observer* the "causes" chronologically preced their own "effects" (for the definition of causes and effects see e.g. Caldirola and Recami[11]). Let us recall that in Sect. 2 the initial statement of the Third Postulate has been shown to be equivalent — as it follows from Postulates 1) and 2 — to the more natural Assumption that ⟨⟨ negative-energy objects travelling forward in time do *not* exist⟩⟩ .

4.1. Existence of an invariant speed

Let us initially *skip* the Third Postulate.

Since 1910 it has been shown (Ignatowski[23], Frank and Rothe[24], Hahn[25], Lalan[26], Severi[27], Agodi[28], Di Jorio[29]) that the postulate of the light-speed invariance is not strictly necessary, in the sense that ourPostulates 1) and 2) *imply* the existence of an invariant speed (*not* of a maximal speed, however). In fact, from the first two Postulates it follows (Rindler[30], Berzi and Gorini[31], Gorini and Zecca[32] and refs. therein, Lugiato and Gorini[33]) that one *and only one* quantity w^2 — having the physical dimensions of the square of a speed — must exist, which has the same value according to

all inertial frames:

$$w^2 = \text{invariant.} \tag{13}$$

If one assumes $w = \infty$, as done in Galilean Relativity, then one would get Galilei-Newton physics; in such a case the invariant speed is the infinite one: $\infty \oplus v = \infty$, where we symbolically indicated by $\oplus$ the operation of speed composition.

If one assumes the invariant speed to be finite and real, then one gets immediately Einstein's Relativity and physics. Experience has actually shown us the speed c of light in vacuum to be the (finite) invariant speed: $c \oplus v = c$. In this case, of course, the infinite speed is no more invariant: $\infty \oplus v = V \neq \infty$. It means that in SR the operation $\oplus$ is not the operation + of arithmetics.

Let us notice once more that the unique role in SR of the light-speed c in vacuum rests on its being invariant and not the maximal one (see e.g. Shankara[34], Recami and Modica[17]); if tachyons — in particular infinite-speed tachyons — exist, they could *not* take over the role of light in SR (i.e. they could not be used by different observers to compare the sizes of their space and time units, etc.), just in the same way as bradyons cannot replace photons. The speed c turns out to be a limiting speed; but any limit can possess *a priori* two sides (Fig. 4).



Fig.4(a)



Fig.4(b)

4.2. The Problem of Lorentz Transformations.

Of course one can substitute the light-speed invariance Postulate for the assumption of space-time homogeneity and space isotropy (see the Second Postulate).

In any case, from the first two Postulates it follows that the transformations connecting two generic inertial frames $\underline{f}$, $\underline{f}'$, a priori with $-\infty < |u| < +\infty$,

$$dx'_\mu = G^\nu_\mu \, dx_\nu \tag{14}$$

must (cf. Sect. 2):

(i) transform inertial motion into inertial motion;

(ii) form a group **G**;

(iii) preserve space isotropy;

(iv) leave the quadratic form invariant, *except for its sign* (Rindler[35], Landau and Lifshitz[36]):

$$dx'_\mu dx'^\mu = \pm \, dx_\mu dx^\mu \ . \tag{15}$$

Notice that eq.(15) imposes — among the others — the light-speed to be invariant (Jammer[37]). Eq.(15) holds for any quantity dx_μ (position, momentum, velocity, acceleration, current, etc.) that be a G-*fourvector*, i.e., that behaves as a fourvector under the transformations belonging to **G**. If we explicitly confine ourselves to slower-than-light relative speeds, $u^2 < c^2$, then we have to skip in eq.(15) the sign minus, and we are left with eq.(2) of Sect.2. In this case, in fact, one can start from the identity transformation $G = \mathbf{1}$, which requires the sign plus, and then retain such a sign for continuity reasons.

On the contrary, the sign minus will play an important role when we are ready to go beyond the light-cone discontinuity. In such a perspective, let us preliminary clarify — on a formal ground — what follows (Maccarrone and Recami[21,22]).

4.3. Orthogonal and Antiorthogonal Transformations: Digression

4.2.1. Let us consider a space having, in a certain initial base, the metric $g^{\mu\nu}$, so that for vectors dx^{α} and tensors $M^{\alpha\beta}$ it is

$$dx^{\alpha} = g^{\alpha\beta}\, dx_{\beta} \quad ; \qquad M^{\alpha\beta} = g^{\alpha\gamma} g^{\beta\delta} M_{\gamma\delta} \, .$$

When passing to another base, one writes

$$dx'^{\mu} = g'^{\mu\nu} dx'_{\nu} \quad ; \qquad M'^{\alpha}{}_{\beta} = g'^{\alpha\gamma} M_{\gamma\beta} \, .$$

In the two bases, the scalar products are defined

$$dx_{\alpha} dx^{\alpha} \equiv dx_{\alpha} g^{\alpha\beta} dx_{\beta} \; ; \quad dx'_{\mu} dx'^{\mu} \equiv dx'_{\mu} g'^{\mu\nu} dx'_{\nu} \, ,$$

respectively.

Let us call A the transformation from the first to the second base, in the sense that

$$dx'^{\mu} = A^{\mu}{}_{\rho}\, dx^{\rho} \, ,$$

that is to say

$$dx^{\mu} = (A^{-1})^{\mu}{}_{\rho}\, dx'^{\rho} \, .$$

Now, if we *impose* that

$$dx_{\alpha} dx^{\alpha} = + dx'_{\mu} dx'^{\mu}, \text{ (assumption)} \tag{16}$$

we get

$$g_{\alpha\beta} = g'_{\mu\nu} A^{\mu}{}_{\alpha} A^{\nu}{}_{\beta} \quad ; \tag{17}$$

however, if we *impose* that

$$dx_\alpha dx^\alpha = -dx'_\mu dx'^\mu \ , \ \text{(assumption)} \qquad (16')$$

we get that

$$g_{\alpha\beta} = -g'_{\mu\nu} A^\mu{}_\alpha A^\nu{}_\beta \ . \qquad (17')$$

4.3.2. Let us consider the case (16)-(17), i.e.

$$dx_\alpha dx^\alpha = dx'_\mu dx'^\mu, \ \text{(assumption)} \qquad (16)$$

and let us look for the properties of transformations A which yield

$$g'_{\mu\nu} = g_{\mu\nu} \ . \qquad \text{(assumption)} \qquad (18)$$

It must be

$$g_{\alpha\beta} = g_{\mu\nu} A^\mu{}_\alpha A^\nu{}_\beta ; \qquad \text{i.e.} \qquad g_{\alpha\beta} = A^\mu{}_\alpha A_{\mu\beta} \ , \qquad (17)$$

wherefrom

$$g^{\gamma\alpha} g_{\alpha\beta} = g^{\gamma\alpha} A^\mu{}_\alpha A_{\mu\beta} = A^{\mu\gamma} A_{\mu\beta} = (A^T)^{\gamma\mu} A_{\mu\beta} \ . \qquad (19)$$

At this point, if we impose that in the initial base

$$g_{\mu\nu} \equiv \eta_{\mu\nu} \ , \qquad \text{(assumption)} \qquad (20)$$

then eq.(19) yields

$$\delta^\gamma{}_\beta = (A^T)^{\gamma\mu} A_{\mu\beta} = g^{\mu k} (A^T)^\gamma{}_k g_{\mu\sigma} A^\sigma{}_\beta = \delta^k{}_\sigma (A^T)^\gamma{}_k A^\sigma{}_\beta = (A^T)^\gamma{}_k A^k{}_\beta \ ,$$

that is to say

$$(A^T)(A) = 1 \ . \qquad (21)$$

4.3.3. Now, in the case (16')-(17'), i.e.

$$dx_\alpha dx^\alpha = -dx'_\mu dx'^\mu \ , \quad \text{(assumption)} \tag{16'}$$

when

$$g_{\alpha\beta} = -g'_{\mu\nu} A^\mu{}_\alpha A^\nu{}_\beta , \tag{17'}$$

let us investigate which are the properties of transformations A that yield

$$g'_{\mu\nu} = - g_{\mu\nu} . \quad \text{(assumption)} \tag{18'}$$

In the particular case, again, when

$$g_{\mu\nu} \equiv \eta_{\mu\nu} \ , \quad \text{(assumption)} \tag{20}$$

it must be

$$g_{\alpha\beta} = - (-g_{\mu\nu}) A^\mu{}_\alpha A^\nu{}_\beta ,$$

i.e. transformations A must still be orthogonal:

$$(A^T)(A) = \mathbb{1} . \tag{21}$$

In conclusion, transformations A when *orthogonal* operate in such a way that

either: (i) $dx_\alpha dx^\alpha = + dx'_\mu dx'^\mu$ and $g'_{\mu\nu} = + \eta_{\mu\nu}$, (22a)

or: (ii) $dx_\alpha dx^\alpha = -dx'_\mu dx'^\mu$ and $g'_{\mu\nu} = - \eta_{\mu\nu}$. (22b)

4.3.4. On the contrary, let us now require that

$$dx_\alpha dx^\alpha = -dx'_\mu dx'^\mu \quad \text{(assumption)} \tag{16'}$$

when

$$g_{\alpha\beta} = -g'_{\mu\nu}\, A^{\mu}{}_{\alpha} A^{\nu}{}_{\beta}, \tag{17'}$$

and simultaneously let us look for the transformations A such that

$$g'_{\mu\nu} = +g_{\mu\nu} \ . \text{ (assumption)} \tag{18}$$

In this case, when in particular assumption (20) holds, $g_{\mu\nu} \equiv \eta_{\mu\nu}$, we get that transformations A must be *anti-orthogonal*:

$$(A^{T})(A) = -\mathbb{1}. \tag{23}$$

4.3.5. The same result (23) is easily obtained when assumptions (16) and (18') hold, together with condition (20).

In conclusion, transformations A when *anti-orthogonal* operate in such a way that

$$\text{either: (i)} \quad dx_{\alpha} dx^{\alpha} = -dx'_{\mu} dx'^{\mu} \quad \text{and} \quad g'_{\mu\nu} = +\eta_{\mu\nu} \ , \tag{24a}$$

$$\text{or:} \quad \text{(ii)} \quad dx_{\alpha} dx^{\alpha} = +dx'_{\mu} dx'^{\mu} \quad \text{and} \quad g'_{\mu\nu} = -\eta_{\mu\nu} \ . \tag{24b}$$

4.3.6. For passing from sub- to Super-luminal frames we shall have(see the following) to adopt *antiorthogonal* transformations. Then, our conclusions (22) and (24) show that we will have to impose a sign-change either in the quadratic form (20'), or in the metric (22'), but *not* — of course — in both: otherwise one would get, as known, an ordinary and not a Superluminal transformation (cf. e.g. Mignani and Recami[38]). We expounded here such considerations, even if elementary, since they arose some misunderstandings (e.g., in Kowalczyński[39]). We choose to assume always (unless differently stated in explicit way):

$$g'_{\mu\nu} = + g_{\mu\nu} \ . \tag{25}$$

Let us add the following comments. One could remember the theorems of Riemannian geometry (theorems so often used in General Relativity),

which state the quadratic form to be positive-definite and the $g_{\mu\nu}$-signature to be invariant, and therefore wonder how it can be possible for our anti-orthogonal transformations to act in a different way. The fact is that the pseudo-Euclidean (Minkowski) space-time is *not* a particular Riemannian manifold, but rather a particular Lorentzian (i.e., pseudo-Riemannian) manifold. The space-time itself of General Relativity (GR) is *pseudo-Riemannian* and not Riemannian (only space is Riemannian in GR): see e.g. Sachs and Wu[40]. In other words, the antiorthogonal transformations do *not* belong to the ordinary group of the so-called "arbitrary" coordinate-transformations *usually* adopted in GR, as outlined e.g. by Møller[41]. This has been overlooked, by authors as e.g. Kowalczyński[39]. However, by introducing suitable scale-invariant coordinates (e.g. dilation-covariant "light-cone coordinates"), both sub- and Super-luminal "Lorentz transformations" can be formally written (Maccarrone *et al.*[42] in such a way to preserve the quadratic form, its sign included (see Sect. 5.8).

Throughout this paper we shall adopt (when convenient) natural units [c = 1].

5. A MODEL-THEORY FOR TACHYONS: AN "EXTENDED RELATIVITY" (ER) IN TWO DIMENSIONS.

Till now we have not taken account of tachyons. Let us finally take them into consideration, starting from a *model-theory*, i.e. from "Extended Relativity" (ER) (Maccarrone and Recami[21,22], Maccarrone *et al.*[42], Barut *et al.*[43], Review I) in two dimensions.

5.1. A Duality Principle.

We got from experience that the invariant speed is $w = c$. Once an inertial frame s_o is chosen, the invariant character of the light-speed allows an exhaustive partition of the set $\{\underline{f}\}$, of all inertial frames $\underline{f}$ (cf. Sec. 4), into the two disjoint, complementary subsets $\{s\}$, $\{S\}$ of the frames having speeds $|u| < c$ and $|U| > c$ relative to s_o, respectively. In the following, for simplicity, we shall consider

ourselves as "the observer s_o". At the present time we neglect the *luminal* frames ($u \equiv U = 0$) as "unphysical". The First Postulate requires frames s and S to be equivalent (for such an extension of the criterion of "equivalence" see Caldirola and Recami[11], Recami[6]), and in particular observers S — if they exist — to have at their disposal the same physical objects (rods, clocks, nucleons, electrons, mesons,...) than observers s. Using the language of multidimensional space-times for future convenience, we can say the first two Postulates to require that even observers S must be able to fill their space (as seen by themselves) with a "lattice-work" of meter-sticks and synchronized clocks (Taylor and Wheeler[44]). It follows that objects must exist which are at rest relatively to S and faster-than-light relatively to frames s; this, together with the fact that luxons ℓ show the same speed to any observers s or S, implies that the objects which are braydons $B(S)$ with respect to a frame S must appear as tachyons $T(s)$ with respect to any frame s, and vice-versa:

$$B(S) = T(S) \; ; \quad T(S) = B(s) \; ; \quad \ell(S) = \ell(s). \tag{26}$$

The statement that the terms B,T,s,S do not have an absolute, but only a *relative* meaning, and eq.(26), constitute the so-called *duality principle* (Olkhovsky and Recami[45], Recami and Mignani[46,47], Mignani *et al.*[48], Antippa[49], Mignani and Recami[50]).

This means that the relative speed of two frames s_1, s_2 (or S_1, S_2) will always be smaller than c; and the relative speed between two frames s, S will be always larger than c. Moreover, the above exhaustive partition is invariant when s_o is made to vary inside $\{s\}$ (or inside $\{S\}$), whilst the subsets $\{s\}$, $\{S\}$ get on the contrary interchanged when we pass from $s_o \in \{s\}$ to a frame $S_o \in \{S\}$.

The main problem is finding out *how* objects that are subluminal w.r.t. (= with respect to) observers S appear to observers s (i.e. to us). It is, therefore, finding out the (Superluminal) Lorentz transformations — if they exist — connecting the observations by S with the observations by s.

5.2. Sub- and Super-Luminal Lorentz Transformations: Preliminaries.

We neglect space-time translations, i.e. consider only *restricted* Lorentz transformations. All frames are supposed to have the same event a their origin. Let us also recall that in the chronotopical space Bs are characterized by time-like, ℓs by light-like, and Ts by space-like world-lines.

The ordinary, subluminal Lorentz transformations (LT) from s_1 to s_2, or from S_1 to S_2, are known to preserve the four-vector type. After Sect. 5.1, on the contrary, it is clear that the "Superluminal Lorentz transformations" (SLT) from s to S, or from S to s, must transform time-like into space-like quantities, and vice-versa. With the assumption (25) it follows that in eq.(15) the *plus* sign has to hold for LT's and the *minus* sign for SLTs:

$$ds'^2 = \pm\, ds^2 \quad ; \quad [\underline{u}^2 \lesseqgtr 1] \tag{15}$$

therefore, in "Extended Relativity" (ER), with the assumption (25), the quadratic form

$$ds^2 \equiv dx_\mu dx^\mu$$

is a *scalar* under LTs, but is a *pseudo-scalar* under SLTs. In the present case, we shall write that LTs are such that

$$dt'^2 - dx'^2 = + (dt^2 - dx^2) \; ; \quad [u^2 < 1] \tag{27$\underline{a}$}$$

while for SLTs it must be

$$dt'^2 - dx'^2 = - (dt^2 - dx^2) \; . \quad [u^2 > 1] \tag{27$\underline{b}$}$$

5.3. Energy-Momentum Space.

Since tachyons are just usual particles w.r.t. their own rest

frames $\underline{f}$, where the $\underline{f}$s are Superluminal w.r.t. us, they will possess *real* rest-masses m_o (Recami and Mignani[46], Leiter[5], Parker[52]). From eq.(27b) applied to the energy-momentum vector p^μ, one derives for free tachyons the ralation

$$E^2 - p_x^2 = - m_o^2 < 0 \quad , \quad [m_o \text{ real}] \tag{28}$$

provided that p^μ is so defined to be a **G**-vector (see the following); so that one has (cf. Figs. 5):



Figs. 5

$$+ m_o^2 > 0 \quad \text{for braydons (time-like case)} \qquad (29a)$$

$$p_\mu p^\mu = 0 \quad \text{for luxons (light-like case)} \qquad (29b)$$

$$- m_o^2 < 0 \quad \text{for tachyons (space-like case)} \qquad (29c)$$

Eqs. (27) - (29) tell us that the roles of space and time and of energy and momentum get interchanged when passing from bradyons to tachyons (See Sect. 5.6). Notice that in the present case (eqs. (29)) it is $\mu = 0,1$. Notice also that tachyons slow down when their energy increases and accelerate when their energy decreases. In particular, divergent energies are needed to slow down the tachyons' speed towards its (lower) limit c. On the contrary, when the tachyons' speed tends to infinity, their energy tends to zero; in ER, therefore, energy can be transmitted only at *finite* velocity. From Figs. 5a,c it is apparent that a braydon may have zero momentum (and minimal energy $m_o c^2$), and a tachyon may have zero energy (and minimal momentum $m_o c$); however Bs cannot exist at zero energy, and tachyons cannot exist at zero momentum (w.r.t. the observers to whom they appear as tachyons!). Incidentally, since *transcedent* (=infinite-speed) tachyons do *not* transport energy but do transport momentum ($m_o c$), they allow getting thr *rigid body* behaviour even in SR (Bilaniuk and Sudarshan[53], Review I, Castorina and Recami[54]). In particular, in elementary particle physics, they might a priori be useful for interpreting in the suitable reference frames the diffractive scatterings, elastic scatterings, etc. (Maccarrone and Recami[55] and refs. therein).

5.4. Generalized Lorentz Transformations (GLT): Preliminaries

Eqs. (27a,b), together with requirements (i)-(iii) of Sect. 4.2, finally imply the LTs to be *orthogonal* and the SLTs to be *anti-orthogonal* (Maccarrone *et al.*[42] and refs. therein):

$$G^T G = +\mathbf{1} \qquad \text{(subluminal case: } u^2 < 1); \tag{30a}$$

$$G^T G = -\mathbf{1} \qquad \text{(Superluminal case: } u^2 > 1), \tag{30b}$$

as anticipated at the end of Sect. 4.3. Both sub- and Super-luminal Lorentz transformations (let us call them "Generalized Lorentz transformations", GLT) result to be unimodular. In the two-dimensional case, however, the SLTs can a priori be special or not; to given them a form coherent with the four-dimensional case (cf. also Sects.5.5, 5.6), one is led to adopt SLTs with negative trace: det $SLT_2 = -1$. In four dimensions, however, all the GLTs will result to be unimodular and special:

$$\det G = +1 , \qquad \forall G \in \mathbf{C} . \tag{31}$$

5.5. The Fundamental Theorem of (Bidimensional) ER.

We have now to write down the SLTs, satisfying the conditions (i)--(iv) of Sect. 4.2 with the sign *minus* in eq.(15), still however with $g'_{\mu\nu} = g_{\mu\nu}$ (cf. Sect.4.3, and Maccarrone and Recami[56]), and show that the GLTs actually from a (new) group $\mathbf{C}$. Let us remind explicitly that an essential ingredient of the present procedure is the assumption that the space-time interval dx^μ is a (chronotopical) vector even with respect to $\mathbf{C}$: see eq. (14).

Any SLT from a sub- to a Super-luminal frame, $s \to S'$, will be identical with a suitable (ordinary) LT — let us call it the "dual" transformation — *except for the fact* that it must change time-like into space-like vectors, and vice-versa, according to eqs.(27b) and (25).

Alternatively, one could say that a SLT is *identical* with its dual subluminal LT, provided that we impose the primed observer S' to use the opposite metric-*signature* $g'_{\mu\nu} = g_{\mu\nu}$, however *without changing* the signs into the definitions of time-like and space-like quantities! (Mignani and Recami[38], Shah[57]).

It follows that a generic SLT, corresponding to a Superluminal

velocity U, will be *formally* expressed by the *product* of the dual LT corresponding to the subluminal velocity $u \equiv 1/U$, *by* the matrix $S \equiv S_2 \equiv \mathbb{1}$, where $\mathbb{1}$ is the two-dimensional identity:

$$\begin{cases} SLT(U) = \pm S \cdot LT(u) & (32) \\ \qquad\qquad [u \equiv \frac{1}{U} \; ; \; u^2 < c^2 \; ; \; U^2 > c^2] & \\ S \equiv i\mathbb{1} \; . & (33) \end{cases}$$

Transformation $S \equiv S_2 \in \mathbb{G}$ plays the role of the "transcendent SLT" since for $u \to 0$ one gets $SLT(U \to \infty) = \pm i\mathbb{1}$. The double sign in eq. (32) is required by condition (ii) of Sect. 4.2; in fact, given a particular subluminal Lorentz transformation L(u) and the $SLT = +iL(u)$, one gets

$$[iL(u)]\,[iL^{-1}(u)] \equiv [iL(u)]\,[iL(-u)] \equiv -\mathbb{1} \; . \qquad (34a)$$

However

$$[iL(u)]\,[-iL^{-1}(u)] \equiv [iL(u)]\,[-iL(-u)] \equiv +\mathbb{1} \; . \qquad (34b)$$

Eqs. (34) show that

$$[iL(u)]^{-1} = -iL^{-1}(u) \equiv -iL(-u) .$$

5.6. Explicit Form of the Superluminal Lorentz Transformations (SLT) in Two Dimensions.

In conclusion, the Superluminal Lorentz transformations $\pm iL(u)$ form a group $\mathbb{G}$ together with *both* the orthochronous *and* the antichronous subluminal LTs of Sect. 2: see Fig. 6. Namely, if $\mathbb{Z}(n)$ is the discrete group of the n-th roots of unity, then the new groups $\mathbb{G}$ of GLTs can be *formally* written down as

$$\begin{cases} \mathbb{G} = \mathbb{Z}(4) \otimes \mathcal{L}_+^\uparrow \; ; & (35) \\ \mathbb{Z}(4) \equiv \{\sqrt[4]{1}\} \equiv \{+1, -1, +i, -i\} \; , & (36) \end{cases}$$



Fig. 6

where $\mathcal{L}_+^\uparrow$ represents *here* the *bidimensional* proper orthochronous Lorentz group. Eq. (35) should be compared with eq.(5'). It is

$$\begin{cases} G \in \mathbf{G} \Longrightarrow -G \in \mathbf{G} \ , \quad \forall G \in \mathbf{G} \ ; \\ \\ G \in \mathbf{G} \Longrightarrow SG \in \mathbf{G} \ , \quad \forall G \in \mathbf{G} \ . \end{cases} \tag{37}$$

The appearance of imaginary units into eqs.(33)-(36) is *only* formal, as it can be guessed from the fact that the transcendent operation $S_2 \equiv \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}$ goes into $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ through a "congruence" (Maccarrone

et al.[42]):

$$\begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} = M \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} M^{T} \ . \tag{38}$$

Actually, the GLTs given by eq.(32)-(33), or (35)-(36), simply represent (Review I, p. 232-233) *all* the space-time pseudo-rotations for $0 \leq \alpha \leq 360^{\circ}$: see Fig. 7. To show this, let us write down explicitly the SLTs in the following formal way

$$\left\{ \begin{array}{l} dt' = \pm\, i\, \dfrac{dt - udx}{\sqrt{1 - u^2}} \ ; \\ \\ dx' = \pm\, i\, \dfrac{dx - udt}{\sqrt{1 - u^2}} \ . \end{array} \right. \quad \left[\begin{array}{c} \text{Superluminal case} \\ u^2 < 1 \end{array} \right] \tag{39}$$

Notice that a priori $\sqrt{\beta^2 - 1} = \pm i\sqrt{1 - \beta^2}$, since $(\pm\, i)^2 = -1$. Moreover, we shall always understand that $\sqrt{1 - \beta^2}$ for $\beta^2 > 1$ represents the upper half-plane solution.

The two-dimensional space-time $M(1,1) \equiv (t,x)$ can be regarded as a complex-plane; so that the imaginary unit

$$i \equiv \exp\left[\tfrac{1}{2}\, i\, \pi\right] \tag{40}$$

operates there as a 90° pseudo-rotation. The same can be said, of course, for the operation S_2; in accord with eq.(38). Moreover, with regard the *axes* $\underline{x}'$, $\underline{t}'$, $\underline{x}$, $\underline{t}$, both observers s_o, S' will agree in the case of a SLT that: $\underline{t}' \equiv \underline{x}$; $\underline{x}' \equiv \underline{t}$. It follows that eqs.(39) can be immediately rewritten:

$$\left\{\begin{array}{l} dt' = \pm \dfrac{dx - udt}{\sqrt{1-u^2}} \ ; \\ \\ dx' = \pm \dfrac{dt - udx}{\sqrt{1-u^2}} \ , \end{array}\right. \qquad \left[\begin{array}{c}\text{Superluminal case} \\ u^2 < 1\end{array}\right] \tag{39'}$$

where the roles of the space and the time coordinates appear interchanged, but the imaginary units *disappeared*.

Let us now take advantage of a very important symmetry property of the ordinary Lorentz boosts, expresse by the *identities*

$$\left\{\begin{array}{l} \dfrac{dx - udt}{\sqrt{1-u^2}} \equiv - \dfrac{dt - Udx}{\sqrt{U^2 - 1}} \ ; \\ \\ \dfrac{dt - udx}{\sqrt{1-u^2}} \equiv - \dfrac{dx - Udt}{\sqrt{U^2 - 1}} \ . \end{array}\right. \qquad [U \equiv 1/u] \tag{41}$$

Eqs.(39') eventually write

$$\left\{\begin{array}{l} dt' = \mp \dfrac{dt - Udx}{\sqrt{U^2 - 1}} \ ; \\ \\ dx' = \mp \dfrac{dx - Udt}{\sqrt{U^2 - 1}} \ , \end{array}\right. \qquad \left[\begin{array}{c}\text{Superluminal case} \\ U^2 > 1\end{array}\right] \tag{39''}$$

which can be assumed as *the canonical form* of the SLTs in two dimensions. Let us observe that eqs.(39') or (39'') yield for the speed of s_o w.r.t. S':

$$x \equiv 0 \implies \frac{dx'}{dt'} = \pm \left(-\frac{1}{u}\right) = \mp U \ , \quad \left[\begin{array}{c} u^2 < 1 \ ; \quad U^2 > 1 \\ \\ U \equiv 1/u \end{array}\right] \tag{42}$$

where u, U are the speeds of the two *dual* frames s, S'. This confirms that eqs.(39), (39") do actually refer to Superluminal relative motion. Even for eqs.(39) one could have derived that the **G**-vectorial velocity $u^\mu \equiv dx^\mu/d\tau_o$ (see the following) changes under trabsformation (39) in such a way that $u'_\mu u'^\mu = - u_\mu u^\mu$; so that from $u_\mu u^\mu = +1$ it follows $u'_\mu u'^\mu = -1$ (that is to say, bradyonic speeds are transformed into tachyonic speeds).

The group **G** of the GLTs in two dimensions can be finally written (Fig. 6):

$$G = \{+L^\uparrow\} \cup \{-L^\uparrow\} \cup \{-SL^\uparrow\} \cup \{+SL^\uparrow\} \ ; \tag{35'}$$

$$S \equiv S_2 \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} . \tag{36'}$$

Notice that the transcendent SLT, S, when applied to the motion of a particle, just interchanges the values of energy and impulse, as well as of time and space: Cf. also Sects.5.2, 5.3 (Review I; see also Vysin[58]).

5.7. Explicity Form of GLTs

The LTs and SLTs together, i.e. the GLTs, can be written of course in a form covariant under the whole group **G**; namely, in "G-covariant" form, they can be written (Figs.7):



Fig.7(a)

$$\begin{cases} dt' = \pm \dfrac{dt - udx}{\sqrt{|1-u^2|}} \\ \\ dx' = \pm \dfrac{dx - udt}{\sqrt{|1-u^2|}} \end{cases} \quad ; \quad \left[\begin{array}{c}\text{Generalized case} \\ -\infty < u < +\infty\end{array}\right] \tag{43}$$

or rather (Recami and Mignani[47]), in terms of the continuous parameter $\sigma \in [0,2\pi]$,



Fig.7(b)

$$\begin{cases} dt' = \Omega\gamma_o(dt - dx\ \mathrm{tg}\,\sigma)\ ; \\ dx' = \Omega\gamma_o(dx - dt\ \mathrm{tg}\,\sigma)\ , \end{cases} \qquad \begin{bmatrix} -\infty < u < +\infty \\ 0 \leq \sigma \leq 2\pi \end{bmatrix} \tag{43a}$$

with

$$\begin{cases} u \equiv \mathrm{tg}\,\sigma\ ;\ \Omega \equiv \Omega\ (\sigma) \equiv \dfrac{\cos\sigma}{|\cos\sigma|}\ \delta^2\ ;\ \delta \equiv +\sqrt{\dfrac{1 - \mathrm{tg}^2\sigma}{|1 - \mathrm{tg}^2\sigma|}}\ ; \\ \gamma_o = +\ (|1 - \mathrm{tg}^2\sigma|)^{-1/2}\ , \end{cases} \tag{43b}$$

where the form (43a) of the GLTs explicitly shows *how* the signs in front of t', x' *succeed* one another as functions of u, or rather of σ (see also the figs. 2-4 and 6 in Review I).

Apart from Somigliana's early paper, only recently rediscovered (Caldirola *et al.*[59]), the eqs. (39"),(43) first appeared in Olkhovsky and Recami[60], Recami and Mignani[46], Mignani *et al.*[48], and then — independently — in a number od subsequent papers: see e.g. Antippa[49] and Ramanujam and Namasivayam[61]. Eqs. (39'), (39") have been shown by Recami and Mignani[46] to be equivalent to the pioneering — even if more complicated — equations by Parker[52]. Only in Mignani *et al.*[48], however, it was first realized that eqs. (39)-(43) need their double sign, necessary in order that *any* GLT admits an inverse transformation (see also Mignani and Recami[50]).

5.8. The GLTs by Discrete Scale Transformations.

If you want, you can regard eqs. (39')-(39") as entailing a "reinterpretation" of eqs. (39), — such a reinterpretation having nothing to do, of course, with the Stückelberg-Feynman "switching procedure", also known as "reinterpretation principle" ("RIP").— Our interpretation procedure, however, not only is straightforward (cf. eqs. (38), (40)), but has been also rendered *automatic* in terms of new, scale-invariant "light-cone coordinates" (Maccarrone *et al.*[42]).

Let us first rewrite the GLTs in a more compact form, by the language of the discrete (real or imaginary) scale transformations (Pavšič and Recami[62], Pavšič[63]):

$$ds'^2 = \rho^2 ds^2 \quad ; \quad \rho^2 = \pm 1 \quad ; \tag{15'}$$

notice that, in eq. (36), $\mathbb{Z}(4)$ is nothing but the discrete group of the dilations $D : x'_\mu = \rho x_\mu$ with $\rho^2 = \pm 1$. Namely, let us introduce the new (discrete) dilation-invariant coordinates (Kastrup[64])

$$\eta^\mu \equiv k x^\mu \quad , \quad [k = \pm 1 \, , \pm i] \tag{44}$$

k being the *intrinsic* scale-factor of the considered object: and let us observe that, under a dilation D, it is $\eta'_\mu = \eta_\mu$ with $\eta'_\mu \equiv k'x'_\mu$, while $k' = \rho^{-1}k$. Braydons (antibraydons) correspond to $k = +1$ ($k=-1$), whilst tachyons and antitachyons correspond to $k = \pm i$. It is interesting that in the present formalism the quadratic form $d\sigma^2 \equiv d\eta_\mu d\eta^\mu$ is invariant, its sign included, under *all* the GLTs:

$$d\sigma'^2 = + d\sigma^2 \quad , \quad \forall G \in \mathbf{G}. \tag{15"}$$

Moreover, under an orthochronous Lorentz transformation $L \in \mathcal{L}^\uparrow_+$, it holds that $\eta'_\mu = L^\mu_{\ \nu}\eta^\nu$; $k' = k$.

It follows —when going back to eq.(14), i.e. to the coordinates x^μ, k— that the generic GLT = G can be written in two dimensions

$$\begin{cases} G = k'^{-1} L k & \left[\rho^2, k^2 = \pm 1\right] \\ L \in \mathcal{L}^\uparrow_+ \ . & \left[k' = \rho^{-1} k\right] \end{cases} \tag{45}$$

5.9. The GLTs in the "Light-Cone Coordinates". *Automatic* Interpretation.

It is known (Bjorken *et al.*[65]) that the ordinary subluminal (proper, orthochrounous) boosts along x can be written in the generic form:

$$\begin{cases} d\xi' = \alpha d\xi \ ; \ d\zeta' = \alpha^{-1} d\zeta \ ; \ dy' = \dfrac{\alpha}{|\alpha|} dy \ ; \ dz' = \dfrac{\alpha}{|\alpha|} dz \ ; \ [0 < \alpha < +\infty] \\ \dfrac{\alpha - \alpha^{-1}}{\alpha + \alpha^{-1}} = u \ ; \ u^2 < 1 \ , \end{cases}$$

in terms of the light-cone coordinates (Fig. 8):

$$\xi \equiv t - x \ ; \ \zeta \equiv t + x' \ ; \ y \ ; \ z. \tag{46}$$



Fig. 8

It is interesting that the orthochronous Lorentz boosts along x just correspond to a dilation of the coordinates ξ,ζ (by the factors α and α^{-1}, respectively, with α any *positive real* number). In particular for $\alpha \to +\infty$ we have $u \to c^-$ and for $\alpha \to 0^+$ we have $u \to -(c^-)$. It is apparent that $\alpha = e^R$, where R is the "rapidity".

The proper *antichronous* Lorentz boosts correspond to the negative real α values (which still yield $u^2 < 1$).

Recalling definitions (44), let us eventually introduce the following *scale-invariant* "light-cone coordinates":

$$\varphi \equiv \eta^{(0)} - \eta^{(1)} \; ; \quad \psi \equiv \eta^{(0)} + \eta^{(1)} \; ; \; \eta^{(2)} \; ; \; \eta^{(3)} \; . \tag{47}$$

In terms of coordinates (47), *all* the two-dimensional GLTs (both sub- and Superluminal) can be expressed in the synthetic form (Maccarrone *et al.*[42]):

$$\begin{cases} d\varphi' = \alpha d\varphi \; ; \quad d\psi' = \alpha^{-1} \, d\psi \; ; \quad [\,|u| \lesseqgtr 1\,] \\ k' = \rho^{-1} k \; ; \quad \alpha \equiv \rho a \; ; \; a \in (0,+\infty) \; ; \; \rho^2 = \pm 1 \; , \end{cases} \tag{48}$$

and all of them preserve the quadratic form, its sign included: $\varphi'\psi' = \varphi\psi$.

The point to be emphasized is that eqs.(48) in the Superluminal case yield directly eq.(39"), i.e. they *automatically* include the "re-interpretation" of eqs.(39). Moreover, eqs.(48) yield

$$\begin{cases} u = \dfrac{\alpha - \alpha^{-1}}{\alpha + \alpha^{-1}} \; ; \\ \qquad\qquad \left[\begin{array}{l} u^2 \lesseqgtr 1 \; ; \\ 0 < a < +\infty \end{array}\right] \\ \alpha = \rho a \; ; \; \rho^2 = \pm 1 \; ; \end{cases} \tag{49}$$

i.e. also in the Superluminal case they forward the correct (faster--than-light) relative speed *without any need of "reinterpretation".*

5.10. An Application

As an application of eqs.(39"), (43), let us consider a tachyon having (real) proper-mass m_o and moving with speed V relatively to us: then we shall observe the relativistic mass

$$m = \frac{m_o}{(|1 - V^2|)^{1/2}} \equiv \frac{-im_o}{(1 - V^2)^{1/2}} \equiv \frac{m_o}{(V^2 - 1)^{1/2}} \; , \; [V^2 > 1 \; ; \; m_o \text{ real}]$$

and, more in general (in G-covariant form):

$$m = \pm \frac{m_o}{(|1 - v^2|)^{1/2}} \quad , \quad [-\infty < v < +\infty] \tag{50}$$

so as anticipated in Fig. 4a.

For other applications, see e.g. Review I; for instance: (i) for the generalized "velocity composition law" in two dimensions see eq. (33) and Table I in Review I; (ii) for the generalization of the phenomenon of Lorentz contraction/dilation see Fig. 8 of Review I.

5.11. Dual Frames (Or Objects)

Eqs.(32) and follows. show that a one-to-one correspondence

$$v \longleftrightarrow \frac{c^2}{v} \tag{51}$$

can be set between subluminal frames (or objects) with speed $v < c$ and Superluminal frames (or objects) with speed $V \equiv c^2/v > c$. In such a particular conformal mapping (*inversion*) the speed c is the "united" one, and the speeds zero, infinite correspond to each other. Cf. also Fig. 9, which illustrates the important equation (32). In fact (Review I) the *relative* speed of two "dual" frames s,S (frames dual one to the other are characterized in Fig. 9 by AB being orthogonal to the u-axis) is *infinite*; the figure geometrically depicts, therefore, the circumstance that $(s_o \to S) = (s_o \to s) \cdot (s \to S)$, i.e. the fundamental theorem of the (bidimensional) "Extended Relativity":((The SLT : $s_o \to S(U)$ is the product of the LT : $s_o \to s(u)$, where $u \equiv 1/U$, by the *transcendent* SLT)) : Cf. Sect. 5.5, eq.(32). (Mignani and Recami[50]).

Even in more dimensions, we shall call "dual" two objects (or frames) moving along the same line with speeds satisfying eq.(51):

$$vV = c^2 \quad , \tag{51'}$$

i.e. with infinite *relative* speed. Let us notice that, if p^μ and P^μ are the energy-momentum vectors of the two objects, then the condition of *infinite* relative speed writes in G-invariant way as

$$p_\mu P^\mu = 0 \qquad (51'')$$



Fig. 9

5.12. The "Switching Principle" for Tachyons.

The problem of the double sign in eq.(50) has been already taken care of in Sect. 2 for the case of braydons (eq.(9)).

Inspection of Fig. 5c shows that, in the case of tachyons, it is enough a (suitable) ordinary subluminal orthochronous Lorentz transformation $L^\uparrow$ to transform a positive-energy tachyon T into

a negative-energy tachyon T'. For simplicity let us here confine ourselves, therefore, to transformations $L \equiv L^{\uparrow} \in \mathcal{L}_{+}^{\uparrow}$, acting on free tachyons.

On the other hand, it is wellknown in SR that the chronological order along a space-like path is *not* $\mathcal{L}_{+}^{\uparrow}$-invariant.

However, in the case of Ts it is even clearer than in the bradyon case that the same transformation L which inverts the energy-sign will also reverse the motion-direction in time (Review I, Recami[66], Caldirola and Recami[67], see also Garuccio *et al.*[2]). In fact, from Fig. 10 we can see that for going from a positive-energy state $\underline{T}_i$ to a negative-energy state $\underline{T}'_f$ it is necessary to bypass the "transcendent" state $\underline{T}_\infty$ (with $V = \infty$). From Fig. 11a we see moreover that, given in the initial frames s_o a tachyon T travelling e.g. along the positive x-axis with speed V_o, the "critical observer" (i.e. the ordinary subluminal observer $s_c \equiv (t_c, x_c)$ seeing T with *infinite* speed) is simply the one whose space-axis x is superimposed to the world-line OT; its speed u_c w.r.t. s_o, along the positive x-axis, is evidently

$$u_c = c^2/V_o \; ; \quad u_c V_o = c^2 \; , \quad [\text{"critical frame"}]$$

dual to the tachyon speed V_o. Finally, from Fig. 10 and Fig. 11b we conclude that any "trans-critical" observer $s' \equiv (t', x')$ such that $u'V_o > c^2$ will see the tachyon T not only endowed with negative energy, but also travelling backwards in time. Notice, incidentally, that nothing of this kind happens when $u V_o < 0$, i.e. when the final frame moves in the direction opposite to the tachyon's.

Therefore Ts display negative energies in the same frames in which they would appear as "going backwards in time", and vice-versa. As a consequence, we can — and must — apply also to tachyons the Stückelberg-Feynman "switching procedure" exploited in Sects. 2.1-2.3. As a result, point $\underline{A}'$ (Fig. 5c) or point $\underline{T}'_f$ (Fig. 10) do *not* refer to a "negative-energy tachyon moving backwards in time", but rather to an *antitachyon* $\overline{T}$ moving the opposite way (in space), forward in time,



Fig. 10



Fig.11(a)

and with positive energy. Let us repeat that the "switching" never comes into the play when the sign of u is opposite to the sign of V_o. (Review I, Recami[68], Caldirola and Recami[11]).

The "Switching Principle" has been first applied to tachyons by Sudarshan and coworkers (Bilaniuk *et al.*[69]; see also Gregory[70]).

Recently Schwartz[71] gave the switching procedure an interesting formalization, in which — in a sense — it becomes "automatic".

5.13. Sources and Detectors. Causality.

After the consideration in the previous Sect.5.12, i.e. when we apply our Third Postulate (Sect.4) also to tachyons, we are left with *no* negative energies (Recami and Mignani[72]) and with *no* motions backwards in time (Maccarrone and Recami[55,73] and refs. therein).

Let us remind, however, that a tachyon T can be transformed into an antitachyon $\overline{T}$ "going the opposite way in space" even by (suitable) ordinary subluminal Lorentz transformation $L \in \mathcal{L}_+^\uparrow$. It is always essential, therefore, when dealing with a tachyon T, to take into proper consideration also its *source* and *detector*, or at least to refer T to an "interaction-region". Precisely, when a tachyon overcomes the divergent speed, it passes from appearing e.g. as a tachyon T *entering* (leaving) a certain interaction-region to appearing as the antitachyon $\overline{T}$ *leaving* (entering) that interaction-region (Arons and Sudarshan[74], Dhar and Sudarshan[74], Glück[75], Baldo *et al.*[76], Camenzind[77]). More in general, the "trans-critical" transformations $\overline{L} \in \mathcal{L}_+^\uparrow$ (cf. the caption of Fig. 11b) lead from a T emitted by A and absorbed by B to its $\overline{T}$ emitted by B and absorbed by A (see Figs.1 and 3b, and Review I).

The already mentioned fact (Sect.2.2) that the Stückelberg - Feynman-Sudarshan "switching" exchanges the roles of source and detector (or, if you want, of "cause" and "effect") led to a series of apparent "causal paradoxes" which — even if easily solvable, at least in microphysics — gave rise to much perplexity in the literature.

We shall deal with the causal problem elsewhere. Let us here anticipate that, — even if in ER the judgement about which is the "cause" and which is the "effect", and even more about the very existence of a "causal connection", is *relative* to the observer —, nevertheless in microphysics the law of "retarded causality" (see our Third Postulate)



Fig.11(b)

remains covariant, since any observers will always see the cause to precede its effect.

Actually, a sensible procedure to introduce Ts in Relativity is postulating *both*: (a) tachyon existence *and* (b) retarded causality, and then trying to build up an ER in which the validity of both postulates is enforced. Till now we have seen that such an attitude —which extends the procedure in Sect.2 to the case of tachyons— has already produced, among the others, the description within Relativity of both matter and antimatter (Ts and $\overline{\text{T}}$s, and Bs and $\overline{\text{B}}$s).

5.14. Braydons and Tachyons. Particles and Antiparticles.

Fig.6 shows, in the energy-momentum space, the existence of *two* different "symmetries", which have nothing to do one with the other.

The symmetry particle/antiparticle is the mirror symmetry w.r.t. the axis $E = 0$ (or, in more dimensions, to the hyperplane $E = 0$).

The symmetry braydon/tachyon is the mirror symmetry w.r.t. the bisectors, i.e. to the two dimensional "light-cone".

In particular, when we confine ourselves to the proper orthochronous subluminal transformations $L^{\uparrow} \in \mathcal{L}_{+}^{\uparrow}$, the "matter" or "antimatter" character is invariant for braydons (but not for tachyons).

We want at this point to put forth explicitly the following simple but important argumentation. Let us consider the two "most-typical" generalized frames: the frame at rest, $s_o \equiv (t,x)$, and its *dual* Superluminal frame (cf.eq.(51) and Fig.9), i.e. the frame $S'_\infty \equiv (t',x')$ endowed with infinite speed w.r.t. s_o. The world-line of S'_∞ will be of course superimposed to the x-axis. With reference to Fig.7b, observer S'_∞ will consider as time-axis t' our x-axis and as space-axis x' our t-axis; and vice-versa for s_o w.r.t. S' . Due to the "extended principle of relativity" (Sect.4), observers s_o, S'_∞ have moreover to be equivalent.

In space-time (Fig.1) we shall have braydons and tachyons going both forward and backwards in time (even if for each observer — e.g. for s_o — the particles travelling into the past have to bear negative energy, as required by our Third Postulate). The observer s_o will of course intepret all —sub- and Superluminal— particles moving *backwards* in *his* time t as antiparticles; and he will be left only with objects going forward in time.

Just the same will be done, in his own frame, by observer S'_∞ since to him all —sub- or Superluminal— particles travelling *backwards* in *his* time t'(i.e. moving along the negative x-direction, according to us) will appear endowed with negative energy. To see this, it is enough to remember that the transcendent transformation S does exchange the values of energy and momentum (cf. eq.(38), the final part of Sect.5.6, and Review I). The same set of braydons and tachyons will be therefore described by S'_∞ in terms of particles and antiparticles all moving along its positive time-axis t'.

But, even if axes t' and x coincide, the observer s_o will see

bradyons and tachyons moving (of course) along both the positive and the negative x-axis! In other words, we have seen the following: The fact S' sees only particles and atiparticles moving along its positive t'-axis *does not mean* at all that s_o sees only braydons and tachyons travelling along his *positive* x-axis! This erroneous belief entered, in connection with tachyons, in the (otherwise interesting) two-dimensional approach by Antippa[49], and later on contibuted to lead Antippa and Everett[78] to violate space-isotropy by conceiving that even in four dimensions tachyons had to move just along a unique, privileged direction — or *"tachyon corridor"*— .

5.15. Totally Inverted Frames.

We have seen that, when a tachyon T appears to overcome the infinite speed (Figs.11a,b), we must apply our Third Postulate, i.e. the "switching procedure". The same holds of course when the considered "object" is a reference frame.

More in general, we can regard the GLTs expressed by eqs. (35')-(36') from the *passive*, and no more from the active, point of view (Recami and Rodrigues[1]). Instead of Fig.6, we get then what depicted in Fig.12. For future convenience, let us use the language of multi-dimensional space-times. It is apparent that the four subsets of GLTs in eq.(35') describe the transitions from the initial frame s_o (e.g. with *right-handed* space-axes) not only to all frames $\underline{f}^R$ moving along x with *all* possible speeds $u = (-\infty,+\infty)$, but also to the "totally inverted" frames $\underline{\tilde{f}}^L = (-1)\underline{f}^R = (\bar{P}\bar{T})\underline{f}^R$, moving as well along x with *all* possible speeds u (cf. Figs.2-6 and 11 in Review I). With reference to Fig. 9, we can say loosely speaking that, if an ideal frame $\underline{f}$ could undergo a whole trip along the axis *(circle)* of the speeds, then —after having overtaken $\underline{f}(\infty) \equiv \underline{f}(U = \infty)$ — it would come back to rest with a *left-handed* set of space-axes and with particles transformed into antiparticles. For further details, see Recami and Rodrigues[1] and refs. therein.



Fig. 12

5.16. About CPT

Let us first remind (sect.5.5) that the product of two SLTs (which is always a *subluminal* LT) can yield a transformation both orthochronous, $L^{\uparrow} \in \mathcal{L}_{+}^{\uparrow}$, and antichronous, $(-\mathbb{1})\cdot L^{\uparrow} \equiv (\overline{P}\,\overline{T})\,L^{\uparrow} = L^{\downarrow} \in \mathcal{L}_{+}^{\downarrow}$ (cf.Sect. 2.3). We can then give eq.(10) the following meaning within ER.

Let us consider in particular (cf.Figs. 7a , b) the antichronous GLT($\theta = 180^{\circ}$) $= -\mathbb{1} \equiv \overline{P}\,\overline{T}$. In order to reach the value $\theta = 180^{\circ}$ starting from $\theta = 0$ we must bypass the case $\theta = 90^{\circ}$ (see Fig.12), where the switching procedure has to be applied (Third Postulate). Therefore:

$$GLT(\theta = 180^{o}) = -\mathbb{1} \equiv \overline{P}\,\overline{T} = CPT \ . \qquad (53)$$

The "total inversion" $-\mathbb{1} \equiv \overline{P}\,\overline{T} = CPT$ is nothing but a particular "rotation" in space-time, and we saw the GLTs to consist in *all* the space-time "rotations" (Sect.5.6). In other words, we can write: $CPT \in \mathcal{G}$, and the CPT-theorem may be regarded as a particular, explicit requirement of SR (as formulated in Sect.2), and *a fortiori* of ER (Mignani and Recami[14,15], and refs. therein, Recami and Ziino[9], Pavšič and Recami[79]). Notice that, in our formalization, the operator CPT is linear and unitary.

5.17. Laws and Descriptions. Interactions and Objects.

Given a certain phenomenon *ph*, the principle of relativity (First Postulate) requires two diferent inertial observers O_1, O_2 to find that *ph* is ruled by the same physical laws, but it does *not* require at all O_1, O_2 to give the same description of *ph* (cf. e.g. Review I; p. 555 in Recami[6]; p. 715, Appendix, in Recami and Rodrigues[1]).

We have already seen in ER that, whilst the "Retarded Causality" is a *law* (corollary of our Third Postulate), the assigment of the "cause" and "effect" labels is *relative* to the observer (Camenzind[77]); and is to be considered a *description-detail* (so as, for instance, the observed colour of an object).

In ER one has to become acquainted with the fact that many description-details, which by chance were Lorentz-invariant in ordinary SR, are no more invariant under the GLTs. For example, what already said (see Sect.2.3, point e)) with regard to the possible non-invariance of the sign of the additive charges under the transformations $L \in \mathcal{L}_+$ holds *a fortiori* under the GLTs, i.e., in ER. Nevertheless, the total charge of an isolated system will have of course to be constant during the time-evolution of the system — i.e. to be conserved — as seen by *any* observer.

Let us refer to the explicit in (Fig.13) (Feinberg[80], Baldo *et al.*[76]), where the pictures (a), (b) are the different descriptions of

the same interaction given by two different (generalized) observers . For instance, (a) and (b) can be regarded as the description, from two *ordinary* subluminal frames O_1, O_2, of one and the same process involving the tachyons a, b (c can be a photon, e.g.). It is apparent that, before the interaction, O_1 sees one tachyons while O_2 see two tachyons. Therefore, the very number of particles —e.g. of tachyons, if we consider only subluminal frames and LTs— observed at a certain time-instant is not Lorentz-invariant. However, the *total* number of particles partecipating in the reaction either in the initial or in the final state *is* Lorentz-invariant (due to our initial three Postulates). In a sense, ER prompts us to deal in physics with interactions rather than with objects (in quantum-mechanical language, with "amplitudes" rather with "states"); (cf. e.g. Gluck[75], Baldo and Recami[81]).

Long ago Baldo *et al.*[76,81] introduced however a vector-space $H = \mathcal{H} \otimes \overline{\mathcal{H}}$ direct product of two vector-spaces $\mathcal{H}$ and $\overline{\mathcal{H}}$, in such a way that any Lorentz transformation was unitary in the H-space even in presence of tachyons. The spaces $\mathcal{H}$ ($\overline{\mathcal{H}}$) were defined as the vector-spaces spanned by the states representing particles and anti-particles only in the initial (final) state.



Figs. 13

5.18. SR with Tachyons in two Dimensions.

Further developments of the classical theory for tachyons in two dimensions, after what precedes, can be easily extracted for example from: Review I and refs. therein; Recami[6,82], Corben[83], Caldirola and Recami[11], Maccarrone and Recami[21,22], Maccarrone *et al.*[42] .

We merely refer here to those papers, and references therein.

Here we shall only make the following (simple, but important) remark. Let us consider two (brayonic) bodies A, B that —owing to mutual attraction— for instance *accelerate while approaching* each other. The situation in Fig.14, where A is chosen as the reference-frame $s \equiv (t,x)$ and, for simplicity, only a discrete change of velocity is depicted. From a Superluminal frame they will be described either as two (anti)tachyons that *accelerate while receding* one from the other [frame $S' \equiv (t',x')$], or as two tachyons that *decelerate while approaching* each other [$S'' \equiv (t'',x'')$]. Therefore, we expect that two tachyons *from the kinematical point of view* will *seem* to suffer a repulsion, if they attract each other in their own rest-frames (and in other frames in which they are subluminal); we shall however see that such a behaviour of tachyons *may* be still considered —from the dynamical, energetical point of view— as due to an *attraction*.

To conclude, let us explicitly remark that the results of the model-theory in two dimensions strongly prompt us to attempt building up a similar theory (based as far as possible on the same Postulates) also in more dimensions.



Fig. 14

6. ACKNOWLEDGEMENTS.

The authors are very grateful, for continuous discussions, to Dr. G. D. Maccarrone and Prof. R. Mignani.

7. REFERENCES.

1) E. Recami and W.A. Rodrigues, *Found. Phys.* 12, 709-718 (1982).

2) A. Garuccio, G.D. Maccarrone, E. Recami and J.P. Vigier, *Lett. Nuovo Cim.* 27, 60-64 (1980).

3) R. Mignani and E. Recami, *Phys. Letters* B65, 148-150 (1976).

4) K. Gödel, in *"A. Einstein: Philosopher-Scientist"* ed. P. A. Schilpp (La Salle, Ill.; Open Court, p. 558), (1973).

5) E. Recami, *Found. Phys.* 8, 329-340 (1978).

6) E. Recami, in *"Albert Einstein* 1879-1979: *Relativity, Quanta and Cosmology"* ed.by F. de Finis and M. Pantaleo (New York: Johnson Rep. Co.), vol. 2, chapt.16, pp. 537-597. This ref. appeared also in Italian: in *"Centenario di Einstein: Astrofisica e Cosmologia, Gravitazione, Quanti e Relatività"* ed. by M. Pantaleo (Florence: Giunti-Barbera, 1979), chapt.18, pp. 1021--1097; and in Russian: in *"Astrofizika, Kvanti i Teorya Otnositelnosti"*, ed. by F.I. Fedorov (Moscow: MIR, pp.53-128), (1982).

7) We shall briefly indicate by "Review I" the review-paper by E. Recami and R. Mignani, *Riv.Nuovo Cim.* 4, 209-290 (1974); (Erratum: p. 398).

8) E. Recami, in *"Tachyons, monopoles, and Related Topics"*, ed. by E. Recami (Amsterdam: North-Holland, p. 3-25) (1978).

9) E. Recami and G. Ziino, *Nuovo Cimento* A33, 205-215 (1976).

10) E.C.G. Stückelberg, *Helv. Phys. Acta* 14, 321, 588 (1971); see also O. Klein, *Zeit. fur Phys.* 53, 157 (1929).

11) P. Caldirola and E. Recami, in *Italian Studies in the Philosophy of Science*, ed. by M. Dalla Chiara (Boston: Reidel, pp. 249-298) (1980).

12) R. Mignani and E. Recami, *Lett.Nuovo Cim.* 18, 5-9 (1977).

13) R. Mignani and E. Recami, *Nuovo Cim.* A24, 438-448 (1974).

14) R. Mignani and E. Recami, *Lett.Nuovo Cim.* 11, 421-426 (1974).

15) R. Mignani and E. Recami, *Int.J.Theor.Phys.* 12,299-320 (1975).

16) M. Pavšič and E. Recami, *Lett.Nuovo Cim.* 34, 357-362 (1982); (Erratum: 35,354).

17) E. Recami and E. Modica, *Lett.Nuovo Cim.* 12, 263-265 (1975).

18) G. Galilei, *"Dialogo...sopra i due massimi sistemi del mondo: Tolemaico e Copernicano"* (Florence: G.B. Landini Pub.) (1632).

19) G. Galilei, *"Dialogue on the Great World Systems: Salusbury Translation"* ed. G. de Santillana (Chicago: Univ. of Chicago Press; p. 199) (1953).

20) V. Gorini, *Comm. Math. Phys.* 21, 150 (1971).

21) G.D. Maccarrone and E. Recami, Report INFN/AE-82/12 (Frascati: INFN, pp.1-39) (1982).

22) G.D. Maccarrone and E. Recami, *Found. Phys.* 14, 367-407 (1984).

23) W.V. Ignatowski, *Phys. Leits.* 21, 972 (1910).

24) P. Frank and H. Rothe, *Ann. der Phys.* 34, 825 (1911).

25) E. Hahn, *Arch.Math. Phys.* 21, 1 (1913).

26) V. Lalan, *Bull.Soc.Math. France* 65, 83 (1937).

27) F. Severi, *"Cinquant'anni di Relatività"* ed. M. Pantaleo (Florence: Giunti-Editrice Universitaria) (1955).

28) A. Agodi, *"Lezioni di Fisica Teorica"* (Catania Univ.,unpublished).

29) M. Di Jorio, *Nuovo Cimento* B22, 70 (1974).

30) W. Rindler, *"Essential Relativity"* (New York : Van Nostrand Reinhold, Sect. 38) (1969).

31) N. Berzi and V. Gorini, *J.Math.Phys.* 10, 1518 (1969).

32) V. Gorini and A. Zecca, *J. Math.Phys.* 11, 2226 (1970).

33) L. Lugiato and V. Gorini, *J.Math.Phys.* 13, 665 (1972).

34) T.S. Shankara, *Found.Phys.* 4, 97 (1974).

35) W. Rindler, *"Special Relativity"* (Edinburg : Oliver & Boyd , p.16) (1966).

36) L. Landau and E. Lifshitz, *"Mecanique"* (Moscow: MIR , 1966) ; *Théorie du Champ"* (1966).

37) M. Jammer, in *"Problems in the Foundations of Physics"*, ed. by G. Toraldo di Francia (1979).

38) R. Mignani and E. Recami,*Lett.Nuovo Cim.* 9, 357-361 (1974).

39) J.K. Kowalczyński, *Int.J. Theor.Phys.* 23, 27 (1984).

40) R. Sachs and W. Wu, *"General Relativity for Mathematicians"* , (Berlin : Springer) (1980).

41) C. Møller, *"The Theory of Relativity"* (Oxford : Oxford Univ. Press , p. 234) (1962).

42) G.D. Maccarrone, M. Pavšič and E. Recami, *Nuovo Cimento*, B73, 91-111 (1983).

43) A.O. Barut, G.D. Maccarrone and E. Recami, *Nuovo Cimento*, A71, 509-533 (1982).

44) E.F. Taylor and J.A. Wheeler, *"Space-Time Physics"* (San Francisco: Freeman) (1966).

45) V.S. Olkhovsky and E. Recami, *Lett. Nuovo Cim.* 1, 165-168 (1971).

46) E. Recami and R. Mignani, *Lett. Nuovo Cim.* 4, 144-152 (1972).

47) E. Recami and R. Magnani, *Lett. Nuovo Cim.* 8, 110-112 (1973).

48) R. Mignani, E. Recami and U. Lombardo, *Lett. Nuovo Cim.* 4, 624--626 (1972).

49) A. Antippa, *Nuovo Cimento* A10, 389-406 (1972).

50) R. Mignani and E. Recami, *Nuovo Cimento* A14, 169-189 (1973); A16, 208 (1973).

51) D. Leiter, *Lett. Nuovo Cim.* 1, 395 (1971).

52) L. Parker, *Phys. Rev.* 138, 2287 (1969).

53) O.M. Bilanink and E.C.G. Sudarshan, *Phys.Today* 22(5),43 (1969).

54) P. Castorina and E. Recami, *Lett. Nuovo Cim.* 22, 195 (1978).

55) G.D. Maccarrone and E. Recami, *Nuovo Cimento* A57, 85-101 (1980).

56) G.D. Maccarrone and E.Recami, *Lett.Nuovo Cim.*4, 251-256 (1982).

57) K.T. Shah, *Lett. Nuovo Cim.* 18, 156 (1977).

58) V. Vyšín, *Nuovo Cimento* A40, 113, 125 (1977).

59) P. Caldirola, G.D. Maccarrone and E. Recami, *Lett. Nuovo Cim.* 29, 241-250 (1980).

60) V.S. Olkhovski and E. Recami, *Vysnik Kiyvskogo Universitetu Ser. Fisiki (Kiev)* 11, 58-64 (1970); *Lett. Nuovo Cim.* 1, 165--168 (1971).

61) G.A. Ramanujam and N. Namasivayan, *Lett. Nuovo Cim.* 6, 245--249 (1973).

62) M. Pavšič and E. Recami, *Lett. Nuovo Cim.* 19, 273-278 (1977).

63) M. Pavšič, in *"Tachyons, monopoles, and Related Topics"*, ed. by E. Recami (Amsterdam: North-Holland, pp. 105-113) (1978).

64) H.A. Kastrup, *Ann. der Phys.* 7, 388 (1962).

65) J.D. Bjorkeen, J.B. Kogut and D.E. Soper, *Phys. Rev.* D3, 1382 (1971).

66) E. Recami, in *"Enciclopedia EST-Mondadori, Annuario'73"* (Milano: Mondadori, pp.85-94) (1973).

67) P. Caldirola and E. Recami, *Epistemologia (Genova)* 1, 263-304 (1978).

68) E. Recami, *Found. Phys.* 8, 329-340 (1978).

69) O.M. Bilanink, V.K. Deshpande and E.C.G. Sudarshan, *Am.J.Phys.* 30, 718 (1962).

70) C. Gregory, *Nature* 192, 1320 (1961); *Phys. Rev.*125,2136 (1962).

71) C. Schwartz, *Phys. Rev.* D25, 356 (1982).

72) E. Recami and R. Mignani, *Lett. Nuovo Cim.* 8, 780-782 (1973).

73) G.D. Maccarrone and E. Recami, *Found. Phys.* 10, 949-957 (1980).

74) M.E. Arons and E.C.G. Sudarshan, *Phys. Rev.* 173, 1622 (1968); J. Dhar and E.C.G. Sudarshan, *Phys. Rev.* 174, 1808 (1968).

75) M. Glück, *Nuovo Cimento* A62, 791 (1969).

76) M. Baldo, G. Fonte and E. Recami, *Lett. Nuovo Cim.* (1st series) 4, 241-247 (1970).

77) M. Camenzind, *Gen. Rel. Grav.* 1, 41 (1970).

78) A.F. Antippa and A.E. Everett, *Phys. Rev.* D8, 2352 (1973).

79) M. Pavšič and E. Recami, *Lett. Nuovo Cim.* 34, 357-367 (1982) ; 35, 354 (1982).

80) G. Feinberg, *Phys. Rev.* 159, 1089 (1967).

81) M. Baldo and E. Recami, *Lett. Nuovo Cim.* (1st series) 2, 643--646 (1969).

82) E. Recami (ed.), *"Tachyons, monopoles, and Related Topics"* (Amsterdam : North-Holland, pp. 1-285) (1982).

83) H.C. Corben, *Nuovo Cimento* A29, 415 (1975); *Int. J. Theor.Phys.* 15, 703 (1976); *Lett. Nuovo Cim.* 22, 116 (1978); and in *"Tachyons, monopoles, and Related Topics"* ed. by E. Recami (Amsderdam: North-Holland, pp. 31-41) (1978).

ISTITUTO NAZIONALE DI FISICA NUCLEARE

IT8500272

Sezione di Catania

INFN/AE-85/2
25 Marzo 1985

E. Recami and W.A.Rodrigues:

A MODEL-THEORY FOR TACHYONS IN TWO DIMENSIONS

Servizio Documentazione
dei Laboratori Nazionali di Frascati

Magnetic Monopoles and Tachyons.

R. MIGNANI and E. RECAMI

Istituto di Fisica Teorica dell'Università - Catania
Istituto Nazionale di Fisica Nucleare - Sezione di Catania
Centro Siciliano di Fisica Nucleare e Struttura della Materia - Catania

(ricevuto il 12 Novembre 1973)

1. – Recently in ref. ([1-3]) the existence has been guessed of a link between magnetic monopoles ([4]) and tachyons ([5,6]). An experimental group ([8]) took seriously that suggestion, and searched for «tachyon monopoles», even if they based their experiment on a theoretical assumption that we deem incorrect ([6,9]).

In the present letter we want preliminarily show the nature of such a connection. We are basing ourselves on special-relativity theory, generalized ([6,7]) for Superluminal inertial frames and faster-than-light objects. For simplicity we shall refer only to x-axis boosts. The main characteristics of «extended relativity» of interest here are:

i) the generalized Lorentz transformations (GLT), which (in the case of x-axis boosts) read ([6,7])

$$(1) \qquad \begin{cases} x = C(x' \cos\varphi + ct' \sin\varphi)\,, & y = (-\eta\delta)y'\,, \\ t = C\left(t' \cos\varphi + \dfrac{x'}{c}\sin\varphi\right), & z = (-\eta\delta)z'\,, \end{cases} \qquad \left[\beta^2 \equiv \left(\frac{u}{c}\right)^2 \gtrless 1\right],$$

(1) L. PARKER: *Phys. Rev.*, **188**, 2287 (1969).
(2) D. WEINGARTEN: Report NBI-HE-71-3, Niels Bohr Institute, Copenhagen (1971), *Ann. of Phys.*, **76**, 510 (1973).
(3) E. RECAMI: in *Enciclopedia EST Mondadori, Annuario 1973* (Milano, 1973), p. 85.
(4) P. A. M. DIRAC: *Proc. Roy. Soc.*, A **133**, 60 (1931); *Phys. Rev.*, **74**, 817 (1948); J. SCHWINGER: *Phys. Rev.*, **144**, 1087 (1966); N. CABIBBO and E. FERRARI: *Nuovo Cimento*, **23**, 1147 (1962). For an extensive bibliography see, *e.g.*, E. AMALDI: in *Old and New Problems in Elementary Particles*, edited by G. PUPPI (New York, 1968); E. AMALDI and N. CABIBBO: in *Aspects of Quantum Theory*, edited by A. SALAM and E. P. WIGNER (Cambridge, 1972).
(5) See, *e.g.*, O. M. P. BILANIUK, V. K. DESHPANDE and E. C. G. SUDARSHAN: *Am. Journ. Phys.*, **30**, 718 (1962).
(6) R. MIGNANI, E. RECAMI: *Riv. Nuovo Cimento* (to appear). See also ref. (3). For an extensive bibliography, see, *e.g.*, ref. (7).
(7) R. MIGNANI and E. RECAMI: *Nuovo Cimento*, **14** A, 169 (1973); **16** A, 208 (1973); E. RECAMI and R. MIGNANI: *Lett. Nuovo Cimento*, **4**, 144 (1972); **8**, 110 (1973); and references therein. See also ref. (4).
(8) D. F. BARTLETT and M. D. LAHANA: *Phys. Rev. D*, **6**, 1817 (1972).
(9) R. MIGNANI and E. RECAMI: *Lett. Nuovo Cimento*, **7**, 388 (1973). See also ref. (6).

where (*)

$$\beta \equiv \operatorname{tg}\varphi \quad [0 \leqslant \varphi \leqslant 2\pi], \quad \gamma \equiv +\,|1-\operatorname{tg}^2\varphi|^{-\frac{1}{2}}, \quad \delta \equiv +\sqrt{\frac{1-\operatorname{tg}^2\varphi}{|1-tg^2\varphi|}},$$

$$\eta \equiv \frac{\cos\varphi}{|\cos\varphi|}\cdot\delta^2, \qquad C \equiv +\sqrt{\frac{1+\operatorname{tg}^2\varphi}{|1-\operatorname{tg}^2\varphi|}};$$

they constitute a new (7) group G;

ii) a « principle of duality » (DP) (6,7), derived from the properties of the group G, which esserts that: « The terms bradyon (7) (B) (usual particles, $v^2 < c^2$), tachyon (T), subluminal inertial frame (s), Superluminal inertial frame (S) do not have an absolute meaning, but only a *relative* one »;

iii) a « rule of tachyonization » (TR) (6,7) (or of the extended relativity), as well derived from the G properties, which states that: « The relativistic laws of mechanics and electromagnetism for tachyons follow by applying a Superluminal Lorentz transformation (SLT)—*e.g.* the *transcendent one* (7) $K_+ \equiv \lim_{\beta\to+\infty} \mathrm{SLT}(\beta>0)$—to the corresponding laws for bradyons ». In fact, particles that are *tachyons* with respect to us (frame s_0) are bradyons with respect to frames S (in particular to *their* own rest frame). And SLT's effect transition from frames S to frames s, and *vice versa*.

2. – The problem of generalizing Maxwell equations for tachyonic sources is not straightforward, since the usual electromagnetic tensor $F_{\mu\nu}$ *a priori* is not expected to be a G-tensor (*i.e.* to behave as a tensor under the extended group G, in particular under SLT's). The problem reduces to generalizing usual Lorentz transformations (10) of the electric field $\boldsymbol{E}$ and of the magnetic field $\boldsymbol{H}$:

$$(2) \qquad \begin{cases} E_x = E'_x, & E_y = \gamma(E'_y + \beta H'_z), & E_z = \gamma(E'_z - \beta H'_y), \\ H_x = H'_x, & H_y = \gamma(H'_y - \beta E'_z), & H_z = \gamma(H'_z + \beta E'_y), \end{cases} \qquad [\beta^2 < 1],$$

for Superluminal velocities. We shall base ourselves (9) on the analogy of « Lorentz transformation » (LT) extension (6,7).

However, let us *first* premise all what follows.

Let us remember that *standard* Maxwell equations read (10)

$$(3) \qquad \partial_\nu F_{\mu\nu} = j_\mu, \qquad \partial_\nu \tilde{F}_{\mu\nu} = 0 \qquad [v^2 < c^2],$$

where $\tilde{F}_{\mu\nu}$ is the *dual* tensor of $F_{\mu\nu}$, and $j_\mu \equiv (\varrho_e, \boldsymbol{j})$ is the *electric* current density four-vector. Notice that we defined (2,11)

$$(4) \qquad \tilde{F}_{\mu\nu} \equiv \tfrac{1}{2}\varepsilon_{\mu\nu\alpha\beta} F_{\alpha\beta} \qquad [\mu, \nu, \alpha, \beta = 0, 1, 2, 3],$$

(*) By the way, notice that in extended relativity both subluminal observers (s) and Superluminal observers (S), being all equivalent (6,7), will see *the same* Minkowsky space, and will use *the same* metric. However, if an observers s (or S) wants to look at space-time not directly, but through the observations of an observer S (or s), *then* he will have to multiply them by i (since what is spacelike for s is timelike for S, and so on). This is the reason why imaginary units enter (only) *the relations between observations from s and from S* (but *not* the « physics » build up by one and the same observer, through his own measures and observations). See R. MIGNANI and E. RECAMI: *Lett. Nuovo Cimento* (to appear).

(10) See, *e.g.*, L. D. LANDAU and E. M. LIFSHITZ: *Theorie du Champ* (Moscow, 1966).

(11) See, *e.g.*, J. M. LEINAAS: *Nuovo Cimento*, **15** A, 740 (1973).

where $\varepsilon_{\mu\nu\alpha\beta}$ is the *real* completely antisymmetric Ricci tensor. It is

$$\tilde{\tilde{F}}_{\mu\nu} = F_{\mu\nu} . \tag{4 bis}$$

Moreover, present «duality» effects the exchanges $\boldsymbol{E} \to i\boldsymbol{H}$, $\boldsymbol{H} \to -i\boldsymbol{E}$.

Let us now introduce the *autodual* (electromagnetic) tensor

$$T_{\mu\nu} \equiv F_{\mu\nu} + \tilde{F}_{\mu\nu} , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} , \tag{5}$$

where $[\mu, \nu = 0, 1, 2, 3]$

$$(T_{\mu\nu}) = \begin{pmatrix} 0 & iE_x - H_x & iE_y - H_y & iE_z - H_z \\ H_x - iE_x & 0 & iE_z - H_z & H_y - iE_y \\ H_y - iE_y & H_z - iE_z & 0 & iE_x - H_x \\ H_z - iE_z & iE_y - H_y & H_x - iE_x & 0 \end{pmatrix} . \tag{6}$$

We are using the metric $(+---)$; however, we write the generic vector as $x \equiv (ct, ix, iy, iz)$, so that $g_{\mu\nu} \equiv \delta_{\mu\nu}$ and we have no distinction between covariant and contravariant components. Summation is understood over the repeated indices (see, *e.g.*, eqs. (3), (4)). Natural units ($c = 1$) will be used when convenient (*).

In terms of $T_{\mu\nu}$, the standard equations (3) can be written together in an essentially single equation:

$$\partial_\nu T_{\mu\nu} = j_\mu , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} \qquad [v^2 < c^2] . \tag{7}$$

As is well known, if magnetic monopoles are supposed to exist, the previous eqs. (3) and (7) may be written [4] in a completely symmetric form, by introduction of the *magnetic*-current density four-vector $g_\mu = (\varrho_m, \boldsymbol{g})$. Equations (3) become

$$\partial_\nu F_{\mu\nu} = j_\mu , \qquad \partial_\nu \tilde{F}_{\mu\nu} = ig_\mu \qquad [v^2 < c^2] , \tag{3'}$$

and eq. (7) then reads

$$\partial_\nu T_{\mu\nu} = j_\mu + ig_\mu , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} \qquad [v^2 < c^2] . \tag{7'}$$

3. – Let us go back to the beginning of Sect. **2**.

It is easy to recognize that the couple E_y, H_z in eqs. (2) behaves as the couple x, t for subluminal LT's. That is to say, for $0 < \beta < 1$, in the plane (E_y, H_z) the axes E'_y, H'_z rotate from 0° to 45°, in anticlockwise and clockwise sense respectively (E'_y starting from E_y, and H'_z from H_z). It is then straightforward to generalize transformations (2) for $1 < \beta < +\infty$, by allowing axes E'_y, H'_z to rotate with continuity *beyond* 45° until 90° (*i.e.*, until when, for $\beta \to +\infty$, E'_y and H'_z coincide with H_z and E_y, respectively). See Fig. 1*a*)-*b*).

(*) As regards the electromagnetic quantities, Heaviside-Lorentz units will be adopted, for practical purposes.



Fig. 1. – Consider an unprimed frame s_0, supposed at rest, and another inertial frame f, to which we shall attribute, every time, all possible collinear velocities u, both subluminal (frames f') and Superluminal (frames f''), along the positive x-axis (*i.e.*, $u \equiv u_x > 0$). Suppose that in f we have a uniform electrostatic field $\boldsymbol{E}$ parallel to the axis y or z (*i.e.*, either $E \equiv E_y$ or $E \equiv E_z$). The cases *a*), *b*) represent how s_0 will see the moving field, according to our eqs. (8*a*), (8*b*). For $0 < u < c$, we shall have the *usual* case *a*), with $0 < \varphi < 45°$. For $0 < u < \infty$, we shall have the case *b*), with $0 < \varphi < 90°$; in particular, for $c < u < \infty$, the angle φ will run from 45° to 90°. We would have met an *analogous* situation when starting from a uniform magnetostatic field, parallel either to the axis y or to the axis z. (See the text.) Notice that here the spaces (E_y, H_z) and (E_z, H_y) are *pseudo-Euclidean*, analogously to space (x, t). On the contrary, *c*) is only a *symbolic graph*, meaning that for $0 < u < c$ we have $E_x(\varphi') = E'_x$, and that for $c < u < \infty$ we have $E_x(\varphi'') = -H_x \equiv iE''_x \equiv H''$. Notice that we are understanding that $\bar{H}_x = E_x$, in Gaussian units. Moreover $(H_0)_x \equiv \bar{H}_x$. Case c) refers for simplicity only to $0 < \varphi < \pi/2$. (See the text.)

Therefore, similarly to eqs. (1), we shall write, for both subluminal *and Superluminal* velocities,

$$\text{(8}a\text{)} \qquad \begin{cases} E_y = C(E'_y \cos\varphi + H'_z \sin\varphi)\,, \\ H_z = C(H'_z \cos\varphi + E'_y \sin\varphi)\,, \end{cases} \qquad [\beta^2 \gtrless 1;\ 0 \leqslant \varphi \leqslant 2\pi].$$

By analogous, «geometrical» extrapolation, we may write, for both subluminal *and Superluminal* velocities,

$$\text{(8}b\text{)} \qquad \begin{cases} E_z = C(E'_z \cos\varphi - H'_y \sin\varphi)\,, \\ H_y = C(H'_y \cos\varphi - E'_z \sin\varphi)\,, \end{cases} \qquad [\beta^2 \gtrless 1].$$

Regarding E_x and H_x, as suggested by eqs. (1) we shall write for Superluminal velocities (*)

$$E_x = iE'_x\,, \qquad H_x = iH'_x \qquad [\beta^2 > 1];$$

in fact, as y, z components behaved as co-ordinates x, t, we assumed now x components to behave as co-ordinate y (or z) in eqs. (1). For both subluminal *and Superluminal* velocities we have (*) (see Fig. 1*c*))

$$\text{(8}c\text{)} \qquad \begin{cases} E_x = E'_x \delta\,, \\ H_x = H'_x \delta\,, \end{cases} \qquad [\beta^2 \gtrless 1].$$

(*) In the following, we shall for simplicity confine ourselves to $-\pi/2 < \varphi < \pi/2$. Notice since now that, due to invariance of our autodual tensor $T_{\mu\nu}$ under «duality», we may consider $iE'_x \equiv -\bar{H}_x$ (where $\bar{H}_x = E_x$, in Gaussian units) and $iH' \equiv \bar{E}_x$ (where $\bar{E}_x = H_x$, in Gaussian units).

Notice explicitly, from eq. (7′), that *multiplication by the imaginary unit i turns quantities due to the electric current into quantities due to the magnetic current* (*). Our transformations (8) may be as well derived by considering the complex vector ([10])

$$\mathscr{F} \equiv \boldsymbol{E} + i\boldsymbol{H}, \qquad \boldsymbol{E} = \operatorname{Re} \mathscr{F}\,, \qquad \boldsymbol{H} = \operatorname{Im} \mathscr{F}\,,$$

and suitable rotations (*).

With transformations (8), we get precisely the interesting result that *under* SLT's *electric and magnetic currents are changed one into the other.* In other words, Superluminal « electric charges » contribute to the electromagnetic field as subluminal magnetic monopoles would do, and *viceversa.* If (bradyonic) monopoles do not exist, then « *electrically charged* » *tachyons will bring into the (generalized) field equation contributions of magnetic-monopole type.*

Namely, let us consider a SLT, *e.g.* the transcendent one (from the unprimed frame to the primed frame) ([6,7]):

$$(1\,bis) \qquad K_+ = \begin{pmatrix} \sigma_2 & 0 \\ 0 & -i\sigma_0 \end{pmatrix} = -(-\sigma_2 \oplus i\sigma_0)\,,$$

where

$$\sigma_0 \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \sigma_2 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

are Pauli's matrices. If, according to DP and TR (Sect. 1), we apply K_+ to eqs. (7), we get (**)

$$(9) \qquad \operatorname{div}\boldsymbol{E}' = 0\,, \qquad \operatorname{div}\boldsymbol{H}' = -\varrho_e' \qquad [\beta^2 > 1]\,,$$

and, by applying K_+ to eqs. (7′),

$$(9') \qquad \operatorname{div}\boldsymbol{E}' = \varrho_m'\,, \qquad \operatorname{div}\boldsymbol{H}' = -\varrho_e' \qquad [\beta^2 > 1]\,.$$

It is easy to verify that, with our eqs. (8), the (Maxwell) field equations (7′) can be written *in G-covariant* form (*i.e.* in a form covariant under the *whole* group G of GLT's)

(*) Therefore, when passing from a Superluminal frame (with speed corresponding to φ_S) to a collinear subluminal frame (with speed corresponding to φ_s), we can better write

$$\begin{cases} E_x(\varphi_s) = iE_x' = iE_x(\varphi_S) = -H_x(\varphi_s)\,, \\ H_x(\varphi_s) = iH_x' = iH_x(\varphi_S) = \;\; E_x(\varphi_s)\,, \end{cases} \qquad [\beta^2 > 1]\,,$$

where, in Gaussian units, $H_x = E_x$ and $E_x = H_x$. *By adopting a different formalism*, one can, for example, write (for $\beta^2 > 1$)

$$\begin{cases} E_x' \xrightarrow{\text{(SLT)}} -H_x\,, \\ H_x' \xrightarrow{\text{(SLT)}} \;\; E_x\,, \end{cases}$$

where now, in Gaussian units, $H_x = E_x'$ and $E_x = H_x'$. These considerations do agree with the well-known invariance of the electromagnetic tensor $T_{\mu\nu}$ under the « duality exchanges ».

(**) By the way, the « transcendent Lorentz transformation » K_+ works so that

$$\begin{cases} E_y' = \;\; H_z\,, & H_y' = -E_z\,, \\ E_z' = -H_y\,, & H_z' = \;\; E_y\,, \end{cases}$$

as *heuristically* forecast in ref. ([1]).

as follows:

$$(10) \qquad \begin{cases} \partial_\nu T_{\mu\nu} = [j_\mu(s) + g_\mu(S)] + i[g_\mu(s) - j_\mu(S)], \\ \tilde{T}_{\mu\nu} \quad = T_{\mu\nu}, \end{cases} \qquad [v^2 \gtrless c^2],$$

where $j_\mu(s)$ and $g_\mu(S)$ are the *subluminal* electric current and the *Superluminal* magnetic current respectively, and so on.

If is assumed magnetic monopoles do not exist, nevertheless (due to tachyon presence) eq. (10) will read

$$(11) \qquad \begin{cases} \partial_\nu T_{\mu\nu} = j_\mu(s) - ij_\mu(S), \\ \tilde{T}_{\mu\nu} \quad = T_{\mu\nu}, \end{cases} \qquad [v^2 \gtrless c^2].$$

Notice that the quantity $T_{\mu\nu}$, which is a tensor under the standard Lorentz group, is no more a usual tensor under the group G, since $T_{\mu\nu}$ under a SLT behaves like a standard tensor, *except*—however—*for* a factor i. That is to say, $T_{\mu\nu}$ is no more supposed to be a standard tensor in eqs. (10)-(11). However, we are assuming $j_\mu \equiv \varrho_0 u_\mu$, where ϱ_0 is the proper charge-density and $u_\mu \equiv \mathrm{d}x_\mu/\mathrm{d}\tau_0$ is the G-fourvelocity ([4]), so that j_μ is a G-fourvector. Let us now for simplicity reduce to subluminat LT's; in this case eqs. (11) are equivalent to the following ones:

$$(12) \qquad \begin{cases} \boldsymbol{\nabla}\cdot\boldsymbol{D} = \varrho(s); \\ \boldsymbol{\nabla}\cdot\boldsymbol{B} = -\varrho(S); \\ \boldsymbol{\nabla}\wedge\boldsymbol{E} = \boldsymbol{j}(S) - \dfrac{\partial \boldsymbol{B}}{\partial t}; \\ \boldsymbol{\nabla}\wedge\boldsymbol{H} = \boldsymbol{j}(s) + \dfrac{\partial \boldsymbol{D}}{\partial t}, \end{cases} \qquad [v^2 \gtrless c^2;\ s = \text{subluminal};\ S = \text{Superluminal}]$$

whose physical meaning is transparent.

All such problems will be dealt with, with more details, in a forthcoming paper, with attention to the experimental consequences too.

Summary. – Connection between magnetic monopoles and faster-than-light particles is shown, in the framework of extended special-relativity theory, by generalizing the Maxwell equations to Superluminal sources.

* * *

Thanks are due to Prof. A. AGODI and Prof. A. GIGLI for kind interest in this work.

MAGNETIC MONOPOLES AND TACHYONS IN SPECIAL RELATIVITY*

E. RECAMI*
Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

and

R. MIGNANI**
International Center for Theoretical Physics, Trieste, Italy

Received 1 December 1975

The *mere* special relativity does not explicitly predict existence of (sub-luminal) monopoles, but on the contrary *explicitly* predicts existence of super-luminal (tachyon) monopoles, with magnetic charge about 100 times less than usually assumed. This fact is relevant also at the light of current experiments looking for magnetic poles.

Since experiments looking for magnetic monopoles failed till now, but a recent claim [1] appeared about a *possible* monopole-detection, it should be interesting to clearly know – and to take possibly into account – the *separate* predictions on the subject of the mere special relativity, on one hand, and of the mere quantum mechanics, on the other hand.

First, we are going to put forth that *special relativity:*

(i) does *not* explicitly predict existence of (slower-than-light) magnetic monopoles;

(ii) *does* explicitly predict, on the contrary, existence of tachyonic (= faster-than-light) "monopoles";

(iii) in its simplest version, predicts the unit magnetic charge for tachyon monopoles to be about one hundred times less than usually [2] assumed ($g = \pm e$, in Gaussian units);

(iv) many good features of the old hypothesis about magnetic monopoles [2] are reproduced, within relativity, by simply taking account of superluminal ($v^2 > c^2$) speeds. In particular, existence of both subluminal ($v^2 < c^2$) and superluminal "electric" charges brings to *fully* symmetrical Maxwell equations [3] (c.f. eqs. (1) in the following).

* Work supported in part by CNR and by INFN-Sezione di Catania.

* On leave from Istituto di Fisica Teorica, Universitá di Catania, Catania, Italy.

** On leave from Istituto di Fisica, Universitá dell'Aquila, L'Aquila, Italy, and from I.N.F.N., Sezione di Roma, Roma.

On the contrary, as regards quantum mechanics, we can say the following. If one assumes *subluminal* magnetic monopole existence, then it seems that the contemporaneous quantization of both electric and magnetic charges follows [2]. This *might* suggest that even subluminal magnetic poles can exist, with unit charge $g = e/2\alpha$, quantity α being the fine-structure constant. Notice, however, that in the present theory the previous one is the *only* argument in favour of subluminal (and large) magnetic charges, since e.g. Maxwell eqs. don't need them to be written in fully symmetrical form.

In fact, let us rebuild [4] a new theory of special relativity *without* assuming *a priori* $|v| < c$. In other words, let us start from the postulates:

1) *Principle of relativity:* laws of mechanics and electro-magnetism are covariant under transitions between two inertial frames, whose relative speed is *a priori* $-\infty < u < +\infty$.

2) *Space is isotropic and space-time homogeneous.*

3) *Principle of retarded causality:* physical signals are transported only by positive-energy objects[‡1].

There follows an "extended relativity" [4], in which light-speed is invariant with respect to *all* inertial frames, both subluminal (s) and superluminal (S), and in which e.g. tachyons do *not* imply any causality violation (as shown in refs. [4, 5] and refs. therein). What is more, the "extended relativity" already proved to be useful even for *usual* particle physics, since *for*

‡1 The usefulness of the third postulate – even in *standard* relativity! – has been shown e.g. in ref. [4].

Magnetic Monopoles and Tachyons.

R. MIGNANI and E. RECAMI

Istituto di Fisica Teorica dell'Università - Catania
Istituto Nazionale di Fisica Nucleare - Sezione di Catania
Centro Siciliano di Fisica Nucleare e Struttura della Materia - Catania

(ricevuto il 12 Novembre 1973)

1. – Recently in ref. ([1-3]) the existence has been guessed of a link between magnetic monopoles ([4]) and tachyons ([5,6]). An experimental group ([8]) took seriously that suggestion, and searched for «tachyon monopoles», even if they based their experiment on a theoretical assumption that we deem incorrect ([6,9]).

In the present letter we want preliminarily show the nature of such a connection. We are basing ourselves on special-relativity theory, generalized ([6,7]) for Superluminal inertial frames and faster-than-light objects. For simplicity we shall refer only to x-axis boosts. The main characteristics of «extended relativity» of interest here are:

i) the generalized Lorentz transformations (GLT), which (in the case of x-axis boosts) read ([6,7])

$$(1) \qquad \begin{cases} x = C(x' \cos\varphi + ct' \sin\varphi)\,, & y = (-\eta\delta)y'\,, \\ t = C\left(t' \cos\varphi + \dfrac{x'}{c}\sin\varphi\right), & z = (-\eta\delta)z'\,, \end{cases} \qquad \left[\beta^2 \equiv \left(\frac{u}{c}\right)^2 \gtrless 1\right],$$

(1) L. PARKER: *Phys. Rev.*, **188**, 2287 (1969).
(2) D. WEINGARTEN: Report NBI-HE-71-3, Niels Bohr Institute, Copenhagen (1971), *Ann. of Phys.*, **76**, 510 (1973).
(3) E. RECAMI: in *Enciclopedia EST Mondadori, Annuario 1973* (Milano, 1973), p. 85.
(4) P. A. M. DIRAC: *Proc. Roy. Soc.*, A **133**, 60 (1931); *Phys. Rev.*, **74**, 817 (1948); J. SCHWINGER: *Phys. Rev.*, **144**, 1087 (1966); N. CABIBBO and E. FERRARI: *Nuovo Cimento*, **23**, 1147 (1962). For an extensive bibliography see, *e.g.*, E. AMALDI: in *Old and New Problems in Elementary Particles*, edited by G. PUPPI (New York, 1968); E. AMALDI and N. CABIBBO: in *Aspects of Quantum Theory*, edited by A. SALAM and E. P. WIGNER (Cambridge, 1972).
(5) See, *e.g.*, O. M. P. BILANIUK, V. K. DESHPANDE and E. C. G. SUDARSHAN: *Am. Journ. Phys.*, **30**, 718 (1962).
(6) R. MIGNANI, E. RECAMI: *Riv. Nuovo Cimento* (to appear). See also ref. (3). For an extensive bibliography, see, *e.g.*, ref. (7).
(7) R. MIGNANI and E. RECAMI: *Nuovo Cimento*, **14** A, 169 (1973); **16** A, 208 (1973); E. RECAMI and R. MIGNANI: *Lett. Nuovo Cimento*, **4**, 144 (1972); **8**, 110 (1973); and references therein. See also ref. (4).
(8) D. F. BARTLETT and M. D. LAHANA: *Phys. Rev. D*, **6**, 1817 (1972).
(9) R. MIGNANI and E. RECAMI: *Lett. Nuovo Cimento*, **7**, 388 (1973). See also ref. (6).

where (*)

$$\beta \equiv \operatorname{tg}\varphi \quad [0 \leqslant \varphi \leqslant 2\pi], \quad \gamma \equiv +\,|1-\operatorname{tg}^2\varphi|^{-\frac{1}{2}}, \quad \delta \equiv +\sqrt{\frac{1-\operatorname{tg}^2\varphi}{|1-tg^2\varphi|}},$$

$$\eta \equiv \frac{\cos\varphi}{|\cos\varphi|}\cdot\delta^2, \qquad C \equiv +\sqrt{\frac{1+\operatorname{tg}^2\varphi}{|1-\operatorname{tg}^2\varphi|}};$$

they constitute a new (7) group G;

ii) a « principle of duality » (DP) (6,7), derived from the properties of the group G, which esserts that: « The terms bradyon (7) (B) (usual particles, $v^2 < c^2$), tachyon (T), subluminal inertial frame (s), Superluminal inertial frame (S) do not have an absolute meaning, but only a *relative* one »;

iii) a « rule of tachyonization » (TR) (6,7) (or of the extended relativity), as well derived from the G properties, which states that: « The relativistic laws of mechanics and electromagnetism for tachyons follow by applying a Superluminal Lorentz transformation (SLT)—*e.g.* the *transcendent one* (7) $K_+ \equiv \lim_{\beta\to+\infty} \mathrm{SLT}(\beta>0)$—to the corresponding laws for bradyons ». In fact, particles that are *tachyons* with respect to us (frame s_0) are bradyons with respect to frames S (in particular to *their* own rest frame). And SLT's effect transition from frames S to frames s, and *vice versa*.

2. – The problem of generalizing Maxwell equations for tachyonic sources is not straightforward, since the usual electromagnetic tensor $F_{\mu\nu}$ *a priori* is not expected to be a G-tensor (*i.e.* to behave as a tensor under the extended group G, in particular under SLT's). The problem reduces to generalizing usual Lorentz transformations (10) of the electric field $\boldsymbol{E}$ and of the magnetic field $\boldsymbol{H}$:

$$(2) \qquad \begin{cases} E_x = E'_x, & E_y = \gamma(E'_y + \beta H'_z), & E_z = \gamma(E'_z - \beta H'_y), \\ H_x = H'_x, & H_y = \gamma(H'_y - \beta E'_z), & H_z = \gamma(H'_z + \beta E'_y), \end{cases} \qquad [\beta^2 < 1],$$

for Superluminal velocities. We shall base ourselves (9) on the analogy of « Lorentz transformation » (LT) extension (6,7).

However, let us *first* premise all what follows.

Let us remember that *standard* Maxwell equations read (10)

$$(3) \qquad \partial_\nu F_{\mu\nu} = j_\mu, \qquad \partial_\nu \tilde{F}_{\mu\nu} = 0 \qquad [v^2 < c^2],$$

where $\tilde{F}_{\mu\nu}$ is the *dual* tensor of $F_{\mu\nu}$, and $j_\mu \equiv (\varrho_e, \boldsymbol{j})$ is the *electric* current density four-vector. Notice that we defined (2,11)

$$(4) \qquad \tilde{F}_{\mu\nu} \equiv \tfrac{1}{2}\varepsilon_{\mu\nu\alpha\beta} F_{\alpha\beta} \qquad [\mu, \nu, \alpha, \beta = 0, 1, 2, 3],$$

(*) By the way, notice that in extended relativity both subluminal observers (s) and Superluminal observers (S), being all equivalent (6,7), will see *the same* Minkowsky space, and will use *the same* metric. However, if an observers s (or S) wants to look at space-time not directly, but through the observations of an observer S (or s), *then* he will have to multiply them by i (since what is spacelike for s is timelike for S, and so on). This is the reason why imaginary units enter (only) *the relations between observations from s and from S* (but *not* the « physics » build up by one and the same observer, through his own measures and observations). See R. MIGNANI and E. RECAMI: *Lett. Nuovo Cimento* (to appear).

(10) See, *e.g.*, L. D. LANDAU and E. M. LIFSHITZ: *Theorie du Champ* (Moscow, 1966).

(11) See, *e.g.*, J. M. LEINAAS: *Nuovo Cimento*, **15** A, 740 (1973).

where $\varepsilon_{\mu\nu\alpha\beta}$ is the *real* completely antisymmetric Ricci tensor. It is

$$\tilde{\tilde{F}}_{\mu\nu} = F_{\mu\nu} . \tag{4 bis}$$

Moreover, present «duality» effects the exchanges $\boldsymbol{E} \to i\boldsymbol{H}$, $\boldsymbol{H} \to -i\boldsymbol{E}$.

Let us now introduce the *autodual* (electromagnetic) tensor

$$T_{\mu\nu} \equiv F_{\mu\nu} + \tilde{F}_{\mu\nu} , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} , \tag{5}$$

where $[\mu, \nu = 0, 1, 2, 3]$

$$(T_{\mu\nu}) = \begin{pmatrix} 0 & iE_x - H_x & iE_y - H_y & iE_z - H_z \\ H_x - iE_x & 0 & iE_z - H_z & H_y - iE_y \\ H_y - iE_y & H_z - iE_z & 0 & iE_x - H_x \\ H_z - iE_z & iE_y - H_y & H_x - iE_x & 0 \end{pmatrix} . \tag{6}$$

We are using the metric $(+---)$; however, we write the generic vector as $x \equiv (ct, ix, iy, iz)$, so that $g_{\mu\nu} \equiv \delta_{\mu\nu}$ and we have no distinction between covariant and contravariant components. Summation is understood over the repeated indices (see, *e.g.*, eqs. (3), (4)). Natural units ($c = 1$) will be used when convenient (*).

In terms of $T_{\mu\nu}$, the standard equations (3) can be written together in an essentially single equation:

$$\partial_\nu T_{\mu\nu} = j_\mu , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} \qquad [v^2 < c^2] . \tag{7}$$

As is well known, if magnetic monopoles are supposed to exist, the previous eqs. (3) and (7) may be written [4] in a completely symmetric form, by introduction of the *magnetic*-current density four-vector $g_\mu = (\varrho_m, \boldsymbol{g})$. Equations (3) become

$$\partial_\nu F_{\mu\nu} = j_\mu , \qquad \partial_\nu \tilde{F}_{\mu\nu} = ig_\mu \qquad [v^2 < c^2] , \tag{3'}$$

and eq. (7) then reads

$$\partial_\nu T_{\mu\nu} = j_\mu + ig_\mu , \qquad \tilde{T}_{\mu\nu} = T_{\mu\nu} \qquad [v^2 < c^2] . \tag{7'}$$

3. – Let us go back to the beginning of Sect. **2**.

It is easy to recognize that the couple E_y, H_z in eqs. (2) behaves as the couple x, t for subluminal LT's. That is to say, for $0 < \beta < 1$, in the plane (E_y, H_z) the axes E'_y, H'_z rotate from 0° to 45°, in anticlockwise and clockwise sense respectively (E'_y starting from E_y, and H'_z from H_z). It is then straightforward to generalize transformations (2) for $1 < \beta < +\infty$, by allowing axes E'_y, H'_z to rotate with continuity *beyond* 45° until 90° (*i.e.*, until when, for $\beta \to +\infty$, E'_y and H'_z coincide with H_z and E_y, respectively). See Fig. 1*a*)-*b*).

(*) As regards the electromagnetic quantities, Heaviside-Lorentz units will be adopted, for practical purposes.



Fig. 1. – Consider an unprimed frame s_0, supposed at rest, and another inertial frame f, to which we shall attribute, every time, all possible collinear velocities u, both subluminal (frames f') and Superluminal (frames f''), along the positive x-axis (*i.e.*, $u \equiv u_x > 0$). Suppose that in f we have a uniform electrostatic field $\boldsymbol{E}$ parallel to the axis y or z (*i.e.*, either $E \equiv E_y$ or $E \equiv E_z$). The cases *a*), *b*) represent how s_0 will see the moving field, according to our eqs. (8*a*), (8*b*). For $0 < u < c$, we shall have the *usual* case *a*), with $0 < \varphi < 45°$. For $0 < u < \infty$, we shall have the case *b*), with $0 < \varphi < 90°$; in particular, for $c < u < \infty$, the angle φ will run from 45° to 90°. We would have met an *analogous* situation when starting from a uniform magnetostatic field, parallel either to the axis y or to the axis z. (See the text.) Notice that here the spaces (E_y, H_z) and (E_z, H_y) are *pseudo-Euclidean*, analogously to space (x, t). On the contrary, *c*) is only a *symbolic graph*, meaning that for $0 < u < c$ we have $E_x(\varphi') = E'_x$, and that for $c < u < \infty$ we have $E_x(\varphi'') = -H_x \equiv iE''_x \equiv H''$. Notice that we are understanding that $\bar{H}_x = E_x$, in Gaussian units. Moreover $(H_0)_x \equiv \bar{H}_x$. Case c) refers for simplicity only to $0 < \varphi < \pi/2$. (See the text.)

Therefore, similarly to eqs. (1), we shall write, for both subluminal *and Superluminal* velocities,

$$\text{(8}a\text{)} \qquad \begin{cases} E_y = C(E'_y \cos\varphi + H'_z \sin\varphi)\,, \\ H_z = C(H'_z \cos\varphi + E'_y \sin\varphi)\,, \end{cases} \qquad [\beta^2 \gtrless 1;\ 0 \leqslant \varphi \leqslant 2\pi].$$

By analogous, «geometrical» extrapolation, we may write, for both subluminal *and Superluminal* velocities,

$$\text{(8}b\text{)} \qquad \begin{cases} E_z = C(E'_z \cos\varphi - H'_y \sin\varphi)\,, \\ H_y = C(H'_y \cos\varphi - E'_z \sin\varphi)\,, \end{cases} \qquad [\beta^2 \gtrless 1].$$

Regarding E_x and H_x, as suggested by eqs. (1) we shall write for Superluminal velocities (*)

$$E_x = iE'_x\,, \qquad H_x = iH'_x \qquad [\beta^2 > 1];$$

in fact, as y, z components behaved as co-ordinates x, t, we assumed now x components to behave as co-ordinate y (or z) in eqs. (1). For both subluminal *and Superluminal* velocities we have (*) (see Fig. 1*c*))

$$\text{(8}c\text{)} \qquad \begin{cases} E_x = E'_x \delta\,, \\ H_x = H'_x \delta\,, \end{cases} \qquad [\beta^2 \gtrless 1].$$

(*) In the following, we shall for simplicity confine ourselves to $-\pi/2 < \varphi < \pi/2$. Notice since now that, due to invariance of our autodual tensor $T_{\mu\nu}$ under «duality», we may consider $iE'_x \equiv -\bar{H}_x$ (where $\bar{H}_x = E_x$, in Gaussian units) and $iH' \equiv \bar{E}_x$ (where $\bar{E}_x = H_x$, in Gaussian units).

Notice explicitly, from eq. (7′), that *multiplication by the imaginary unit i turns quantities due to the electric current into quantities due to the magnetic current* (*). Our transformations (8) may be as well derived by considering the complex vector ([10])

$$\mathscr{F} \equiv \boldsymbol{E} + i\boldsymbol{H}, \qquad \boldsymbol{E} = \operatorname{Re} \mathscr{F}\,, \qquad \boldsymbol{H} = \operatorname{Im} \mathscr{F}\,,$$

and suitable rotations (*).

With transformations (8), we get precisely the interesting result that *under* SLT's *electric and magnetic currents are changed one into the other.* In other words, Superluminal « electric charges » contribute to the electromagnetic field as subluminal magnetic monopoles would do, and *viceversa.* If (bradyonic) monopoles do not exist, then « *electrically charged* » *tachyons will bring into the (generalized) field equation contributions of magnetic-monopole type.*

Namely, let us consider a SLT, *e.g.* the transcendent one (from the unprimed frame to the primed frame) ([6,7]):

$$(1\,bis) \qquad K_+ = \begin{pmatrix} \sigma_2 & 0 \\ 0 & -i\sigma_0 \end{pmatrix} = -(-\sigma_2 \oplus i\sigma_0)\,,$$

where

$$\sigma_0 \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \sigma_2 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

are Pauli's matrices. If, according to DP and TR (Sect. 1), we apply K_+ to eqs. (7), we get (**)

$$(9) \qquad \operatorname{div}\boldsymbol{E}' = 0\,, \qquad \operatorname{div}\boldsymbol{H}' = -\varrho_e' \qquad [\beta^2 > 1]\,,$$

and, by applying K_+ to eqs. (7′),

$$(9') \qquad \operatorname{div}\boldsymbol{E}' = \varrho_m'\,, \qquad \operatorname{div}\boldsymbol{H}' = -\varrho_e' \qquad [\beta^2 > 1]\,.$$

It is easy to verify that, with our eqs. (8), the (Maxwell) field equations (7′) can be written *in G-covariant* form (*i.e.* in a form covariant under the *whole* group G of GLT's)

(*) Therefore, when passing from a Superluminal frame (with speed corresponding to φ_S) to a collinear subluminal frame (with speed corresponding to φ_s), we can better write

$$\begin{cases} E_x(\varphi_s) = iE_x' = iE_x(\varphi_S) = -H_x(\varphi_s)\,, \\ H_x(\varphi_s) = iH_x' = iH_x(\varphi_S) = \;\; E_x(\varphi_s)\,, \end{cases} \qquad [\beta^2 > 1]\,,$$

where, in Gaussian units, $H_x = E_x$ and $E_x = H_x$. *By adopting a different formalism*, one can, for example, write (for $\beta^2 > 1$)

$$\begin{cases} E_x' \xrightarrow{\text{(SLT)}} -H_x\,, \\ H_x' \xrightarrow{\text{(SLT)}} \;\; E_x\,, \end{cases}$$

where now, in Gaussian units, $H_x = E_x'$ and $E_x = H_x'$. These considerations do agree with the well-known invariance of the electromagnetic tensor $T_{\mu\nu}$ under the « duality exchanges ».

(**) By the way, the « transcendent Lorentz transformation » K_+ works so that

$$\begin{cases} E_y' = \;\; H_z\,, & H_y' = -E_z\,, \\ E_z' = -H_y\,, & H_z' = \;\; E_y\,, \end{cases}$$

as *heuristically* forecast in ref. ([1]).

as follows:

$$(10) \qquad \begin{cases} \partial_\nu T_{\mu\nu} = [j_\mu(s) + g_\mu(S)] + i[g_\mu(s) - j_\mu(S)], \\ \tilde{T}_{\mu\nu} \quad = T_{\mu\nu}, \end{cases} \qquad [v^2 \gtrless c^2],$$

where $j_\mu(s)$ and $g_\mu(S)$ are the *subluminal* electric current and the *Superluminal* magnetic current respectively, and so on.

If is assumed magnetic monopoles do not exist, nevertheless (due to tachyon presence) eq. (10) will read

$$(11) \qquad \begin{cases} \partial_\nu T_{\mu\nu} = j_\mu(s) - ij_\mu(S), \\ \tilde{T}_{\mu\nu} \quad = T_{\mu\nu}, \end{cases} \qquad [v^2 \gtrless c^2].$$

Notice that the quantity $T_{\mu\nu}$, which is a tensor under the standard Lorentz group, is no more a usual tensor under the group G, since $T_{\mu\nu}$ under a SLT behaves like a standard tensor, *except*—however—*for* a factor i. That is to say, $T_{\mu\nu}$ is no more supposed to be a standard tensor in eqs. (10)-(11). However, we are assuming $j_\mu \equiv \varrho_0 u_\mu$, where ϱ_0 is the proper charge-density and $u_\mu \equiv \mathrm{d}x_\mu/\mathrm{d}\tau_0$ is the G-fourvelocity ([4]), so that j_μ is a G-fourvector. Let us now for simplicity reduce to subluminat LT's; in this case eqs. (11) are equivalent to the following ones:

$$(12) \qquad \begin{cases} \boldsymbol{\nabla}\cdot\boldsymbol{D} = \varrho(s); \\ \boldsymbol{\nabla}\cdot\boldsymbol{B} = -\varrho(S); \\ \boldsymbol{\nabla}\wedge\boldsymbol{E} = \boldsymbol{j}(S) - \dfrac{\partial \boldsymbol{B}}{\partial t}; \\ \boldsymbol{\nabla}\wedge\boldsymbol{H} = \boldsymbol{j}(s) + \dfrac{\partial \boldsymbol{D}}{\partial t}, \end{cases} \qquad [v^2 \gtrless c^2;\ s = \text{subluminal};\ S = \text{Superluminal}]$$

whose physical meaning is transparent.

All such problems will be dealt with, with more details, in a forthcoming paper, with attention to the experimental consequences too.

Summary. – Connection between magnetic monopoles and faster-than-light particles is shown, in the framework of extended special-relativity theory, by generalizing the Maxwell equations to Superluminal sources.

* * *

Thanks are due to Prof. A. AGODI and Prof. A. GIGLI for kind interest in this work.

Физ.Лэт. т.62B, №1 10 Май 1976

МАГНИТНЫЕ МОНОПОЛИ И ТАХИОНЫ В СПЕЦИАЛЬНОЙ ТЕОРИИ ОТНОСИТЕЛЬНОСТИ.

Э. Реками

Институт Нильса Бора, Университет Копенгагена, Дания

Р. Мигнани

Международный Центр Теоретической Физики, Триест, Италия

Получена 1 Декабря 1975

Сама по себе специальная теория относительности в действительности не предсказывает явно существование (досветовых) монополей, но в противоположность этому явно предсказывает существование сверхсветовых (тахионных) монополей, с магнитными зарядами примерно в 100 раз меньшими, чем обычно предполагается. Этот факт уместен также в свете нынешних экспериментов по поиску магнитных полюсов.

Так как эксперименты по поиску магнитных монополей до сих пор оканчивались неудачей, за исключением недавнего утверждения [1], появившегося *о возможном* детектировании монополей, было бы интересным чётко знать и принять, возможно, во внимание- *различные* предсказания на эту тему самой по себе чистой специальной теории относительности с одной стороны, и самой по себе чистой квантовой механики, с другой стороны.

Во-первых, мы собираемся использовать тот факт, что специальная теория относительности:

1. на самом деле *не* предсказывает явно существования (медленнее –чем -свет) магнитных монополей;

2. *действительно* явно предсказывает, наоборот, существование тахионных (=быстрее –чем- свет) “монополей”;

3. в своей простейшей версии , предсказывает, что единичный магнитный заряд для тахионных монополей примерно в сотню раз меньше, чем обычно [2] преполагается(g=+/-e, в гауссовых единицах);

4. много хороших свойств старых гипотез о магнитных монополях [2] воспроизводится , в рамках теории относительности, простым учётом сверхсветовых (v^2>c^2) скоростей. В частности, существование обоих, как досветовых (v^2<c^2), так и сверхсветовых “электрических” зарядов приводит к *полностью* симметричным уравнениям Максвелла [3] (смотри ур-ние (1) ниже).

Наоборот, что касается квантовой механики, мы можем сказать следующее. Если предположить существование досветовых магнитных монополей , то, по-видимому, одновременное квантование обоих, как элетрического, так и магнитного зарядов следует [2]. Это *может* означать, что даже досветовые магнитные полюса могут существовать, с единичным зарядом $g=e/2\alpha$, где величина α есть постоянная тонкой структуры. Отметим, однако, что в насточщей теории предыдущее обстоятельство не является *единственным* аргументом в пользу досветовых (и больших) магнитных зарядов, раз уж уравнения Максвелла не нуждаются в них для того, чтобы быть записанными в полностью симметричном виде.

Действительно, давайте заново построим новую теорию [4] специальной относительности *без* предположения *априори* |v|<c.Другими словами, давайте стартовать со следующих постулатов:

1)*Принцип относительности:* законы механики и электромагнетизма ковариантны при преобразованиях между двумя инерциальными системами отсчёта, чьи относительные скорости *априори* $-\infty < u < +\infty$.

(2)*Пространство изотропно и пространство-время однородно*.

(3)*Принцип запаздывающей причинности:* физические сигналы переносятся только объектами с положительной энергией.*(Полезность третьего постулата – даже в *стандартной* относитеельности!- была показана , например, в [4])

Отсюда следует “ расширенная относительность”[4] , в которой скорость света инвариантна по отношению ко *всем* инерциальным системам, как досветовым (s) . так и сверхсветовым (S), и в которой , например, тахионы *не* означают какого-либо нарушения причинности (как показано в [4,5], и ссылках там). И даже больше, “расширенная относительность” уже оказалась полезной даже для *обычной* физики частиц, так как *например* позволила вывести “ перекрёстные соотношения” для релятивистских реакций [6].

Момент, который мы хотим особо выделить здесь, это следующее. Если мы рассмотрим существование *электрических* зарядов как досветовых, так и сверхсветовых, (с 4-векторной плотностью тока $J\mu\ (s) \equiv (\rho\ (s), \boldsymbol{J}(s))$ и сверхсветовых (чья 4-векторная плотность тока есть $J\mu\ (S) \equiv (\rho\ (S), \boldsymbol{J}(S))$), то тогда (обобщённые) уравнения Максвелла , как уже было показано в [3,4] ,выглядят, для v^2F c^2,как

div$\boldsymbol{D}$=ρ (s); [для v^2F c^2;

div$\boldsymbol{B}$=-ρ (S); s↔ |v|<c; (1)

rot$\boldsymbol{E}$=$\boldsymbol{J}$(S)-∂ $\boldsymbol{J}/\partial$t; S↔ |v|>c.]

rot$\boldsymbol{H}$=$\boldsymbol{J}$(S)+ ∂ $\boldsymbol{D}/\partial$t.

То есть движущиеся быстрее – чем – свет электрические заряды по предсказаниям (расширенной) теории относительности ведут себя *как* магнитные монополи по имеющимся предположениям должны были бы себя вести (не говоря уже об отличающейся скорости)*(см. также рис.1 в [3]). А именно, сверхсветовой электрический , *положительный* заряд (например,со скоростью V>c вдоль *x*-оси) будет вносить вклад в в полевые уравнения такой, как это как это ожидалось от магнитного *южного* полюса (с [4] v=c^2/V вдоль оси *x*) [4]*(Таким образом, “тахионные электроны” будут появляться с северными магнитным зарядом (+g),а “тахионные протоны” с южным магнитным зарядом (-g), и так далее.) Следовательно, сверхсветовой единичный электрический заряд *e* будет являться нам как (тахионный) “монополь” с единичным магнитным зарядом:

$g=-e$, (в гауссовой системе единиц) (2)

Отсюда следует, что относительность явно предсказывает (тахионный) магнитный минимальный заряд примерно в 100 раз меньше, чем обычно [2] предполается. Другими словами, сама по себе одна лишь расширенная теория относительности явно предсказывает *один* заряд (давайте назовём его “электромагнитным зарядом”), который ведёт себя как “электрический” когда он медленнее, чем, и “магнитный”, когда он быстрее, чем свет*

И *уравнения Максвелла могут быть записаны в полностью симметричной форме*, ур-ние (1), *даже без предположения о существовании (досветовых) монополей.* Как бы то ни было, если досветовые монополи существали,(как *возможно* требуется квантовой механикой),с единичным магнитным зарядом $g=e/2\alpha$ (или $g=e/\alpha$) ,тогда при движении со скоростью больше скорости света они будут возникать как большие (тахионные) электрические заряды $e/2\alpha$ (или e/α).

Более того, давайте заметим, что *универсальность* электромагнитных взаимодействий естественно присутствует в расширенной относительности, раз уж $|g|=|e|$, т.е. только одна константа связи естественно возникает в нашей конструкции, *без какого либо обращения за помощью к квантовой механике,* как и требуется истинным духом самой по себе теории относительности. Наоборот, существование досветовых (большой заряд) магнитных монополей будет нарушать в *специальной теории относительности* универсальность электромагнитного взаимодействия, которая восстанавливается только соотношением Дирака $eg=n\hbar$. В действительности, это соотношение , как было показано, *невыводимо* из классической (= не -квантовой) физики [7].

В соответствии с одной только (расширенной) теорией относительности, все экспериментальные поиски магнитных монополей должны быть переделаны тем, кто действительно ищет “тахионные монополи”*(Действительно, Бартлетт и Лахана [8] уже искали “тахионные монополи”, но безнадёжно, потому что на базе теоретических предположений которые, как мы показали, некорректны: ср.с [9]. Работа до сих пор в состоянии дальнейшего прогресса для достижения лучшего понимания экспериментального поведения тахионных монополей.)т.е. принимая во внимание новую недавно предложенную кинематику (скоростей, больших, чем скорость света) [10] и много меньшую величину соответствующей силы магнитного поля.

Кроме того, экспериментальные планы должны принимать во внимание наше доказательство [11], что в расширенной относительности тахионов *не* ожидается испускание черенковского излучения в вакууме.

Наконец, для того, чтобы поддержать наше уравнение (2), позвольте нам подчеркнуть, что (!) соотношение Дирака $eg=n\hbar/2$ приходит в теорию только когда магнитные монополи предполагаются *досветовыми*; (!!) наоборот, если магнитные монополи рассматриваются как являющиеся *сверхсветовыми*, то тогда расширенная относительность даёт [12] *(Из требования, чтобы инерциальные системы (обе, как s, так и S) были *все* эквивалентны, следует что также и системы S наблюдают электроны, протоны и т.д. . Такие объекты будут появляться в системах s как несущие магнитный заряд $g=+-e$ как получается при применении сверхсветовых преобразованиях Лоренца [4].) альтернативное соотношение $g=ne$. В частности, про “тахионные монополи” можно предсказать, что они испытывают в электромагнитных полях “Силу Лоренца” $\boldsymbol{F=-e(H-V\times E)}$, где $V^2 > c^2$ и гауссова система единиц была использована. Подробности о том, что предстоит, появятся в другом месте [13] Давайте здесь кратко покажем пункт (!), ссылаясь на процедуру, которая следует в [14] .

Кабибо и Феррари [14] , кроме стандартного электромагнитного 4-векторного потенциала $A\mu$, ввели (в дираковский аддитивный член) второй 4-векторный потенциал $B\mu$, когда присутствие магнитных монополей-конечно, предполагающихся ими как досветовых – и использовали метод Мандельштамта [14] для квантования их теории. В этом подходе полевые величины, описывающие электрические заряды (во взаимодействии с электромагнитным полем), есть :

$$\Phi(x;P)=\varphi(x)\exp[-ie\int_{(P)}A\mu(\xi)d\xi\mu],$$

где интеграл оценивается на пространственно-подобном пути P, заканчивающемся в точке x, и где Φ не зависящая от калибровки, выбранной для $A\mu$, но зависящей от пути P :

$$\Phi(x;P')=\Phi(x;P)\exp[-ie/2\int_S F\mu\nu\, d\sigma\,\mu\nu], \quad (3)$$

Причём величина S является поверхностью, ограниченной путями P и $P^{!}$. В этом пункте , если мы предположим урвнение (3) в качестве определения и прмем в рассмотрение также и *досветовые* магнитные полюса, так что $F\mu\ \nu = A\nu / \mu - A\mu / \nu$ - iε μ ν ρ σ $B\sigma / \rho$, то тогда нам придётся наложить дополнительное условие

$$\exp[-ie/2\int_S F\mu\ \nu\, d\sigma\ \mu\ \nu\ ]=1 \quad (4)$$

для того, чтобы это определение (3) действительно *не* зависело от выбранной поверхности S . Условие (4) даёт [14] в точности Дираковское соотношение $eg=nh\#/2$.

Наоборот, при рассмотрении “тахионных монополей”, следует действовать следующим образом. Тахионные полюса в системах S есть ничто иное, как досветовые электрические заряды, и их поля в них описываются стандартным 4-векторным потенциалом $A'\mu$. Затем мы берём 4-векторный потенциал $B\mu$, ассоциированный со сверхсветовыми “магнитными полюсами” *только* как потенциал, трансформированный [12] из $A'\mu$

Подходящим сверхсветовым преобразованием Лоренца [4]L :

$$B\mu = -iL\mu\ \rho\ A^{!\rho} .$$

Так как все соотношения (5-12) ссылки [14] конечно остаются в силе для $A'\mu$ также, как и для

$A\mu$, то для *тахионных монополей* мы в действительности не нуждаемся в дополнительных условиях типа (4), и мы остаёмся только с нашим уравнением (2).

Из последовавшего анализа , даже о возможной роли сверхсветовых скоростей в адронной структуре может быть сделано заключение *(Согласно нам, кварки- это структуры, сделанные из партонов. И кварки могут быть квантованными струнами или петлями из более быстрых, чем свет лептонов (напр. *электронов*), в то время как партоны могут быть этими сверхсветовыми лептонами: см. [12]).

Авторы благодарны Проф. П.Ольсену, А.Агоди, Э. Амальди, Л.Р. Балдини, А.О.Баруту, В.Де Саббата, Е. Феррари, Л. Фонда, Г. Гиакомели, Х.Б. Нильсену, М. Павсичу, А. Саламу, Л.Скарзи, А. Зичичи, и Г. Зиино за дискуссии или их интерес. Один из нас(Э.Р.) благодарит также Проф. О. Бора за проявленное гостеприимство в Институте Нильса Бора.

ЛИТЕРАТУРА

1. P.B.Price, E.K. Shrik, W.Z.Osborne and L.S.Pinsky, Phys.Rev.Lett.35(1975)487.
2. P.A.M.Dirac, Proc.Roy.Soc.A133(1931)60; Phys.Rev.74(1948)817; see also J.Schwinger, Phys.Rev.144(1966)1087.
3. R.Mignani and E.Recami, Lett.Nuovo Cim.9(1974)367.
4. R.Mignani and E.Recami, Rivista N.Cim. 4(1974)209.
5. E.Recami, in Annuario 73, Encyclopedia EST-Mondado (Milano,1973) p.85, and references therein; see e.g. E.C.G. Sudarshan, in Simposia on Theor. Phys. and Math. vol.10 (New York, 1970) p.129; R.G.Root and J.S.Trefil, Lett.Nuovo Cim. 3 (1970)41; J.A.Parmentola and D.D.H. Yee, Phys.Rev.D4(1971)1912.
6. R.Mignani and E.Recami, Int. J. Theor. Phys.12(1975)299; Nuovo Cimento A24 (1974)438; Lett.Nuovo Cim.11(1974)421.
7. H.A.Cohen, Found. Phys. 4(1974) 115.
8. D.F.Bartlett and M.D. Lahana, Phys. Rev.D6 (1972) 1817.
9. R.Mignani and E.Recami, Lett.Nuovo Cim.9(1974)362.
10. R.Mignani and E.Recami, in preparation.
11. R.Mignani and E.Recami, Lett.Nuovo Cim.7(1973)388; Lett.Nuovo Cim.9(1974) 362; Nuovo Cimento B21(1974)210.
12. R.Mignani and E.Recami, preprint IC/75/88 (Trieste,1975), to appear in Nuovo Cimento A.
13. R.Mignani and E.Recami, to be submitted to Phys.Rev.
14. N.Cabibbo and E.Ferrari, Nuovo Cimento 23 (1962) 1147; see also S. Mandelstam, Ann. of Phys. 19(1962) 1.